

A Note on Extensions of Asymptotic Density

A. Blass, R. Frankiewicz, G. Plebanek, C. Ryll–Nardzewski

Abstract

By a density we mean any extension of the asymptotic density to a finitely additive measure defined on all sets of natural numbers. We consider densities associated to ultrafilters on ω and investigate two additivity properties of such densities. In particular, we show that there is a density ν for which $L_1(\nu)$ is complete.

1. Introduction.

We denote the set of natural numbers by ω ($= \{0, 1, 2, \dots\}$), and often regard any $n \in \omega$ as the set $\{0, 1, \dots, n-1\}$. The symbol $\mathcal{P}(\omega)$ stands for the family of all subsets of ω . Recall that the asymptotic density of a set $A \subseteq \omega$, denoted here by $d(A)$, is defined as

$$d(A) = \lim_{n \rightarrow \infty} \frac{|A \cap n|}{n},$$

provided this limit exists. Note that, while d is finitely additive, the domain of d is not an algebra. Several authors have considered extensions of asymptotic density to a finitely additive measure defined on a certain algebra of sets; see for instance Buck [2], Maharam [6], and Mekler [7]. In the sequel, any finitely additive ν defined on $\mathcal{P}(\omega)$ and extending d will be called a *density*.

Following Buck [2] and Mekler [7], we shall consider a certain additivity property of densities. Say that a density ν has property **AP**(*) if for every increasing sequence $(A_i)_{i \geq 1} \subseteq \mathcal{P}(\omega)$ there is a set $B \in \mathcal{P}(\omega)$ such that

- (i) $A_i \subseteq^* B$ for every i , and
- (ii) $\nu(B) = \lim_i \nu(A_i)$.

2000 *Mathematics Subject Classification*: Primary 28A12; Secondary 03E05, 03E35, 11B05.

The first-named author was partially supported by NSF grant DMS-9505118
The other authors were partially supported by KBN grant 2P03A 018 13.

Here and below, we write $A \subseteq^* B$ to denote that the set $A \setminus B$ is finite; in this case we also say that A is *almost included* in B .

Suppose now that a σ -algebra \mathcal{F} of subsets of an arbitrary space X is given, and that ν is a *finitely additive* finite measure defined on \mathcal{F} . One can consider the following natural weakening of **AP**(*): Say that ν has property **AP**(null) if for every increasing sequence $(A_i)_{i \geq 1} \subseteq \mathcal{F}$ there is a set $B \in \mathcal{F}$ such that

- (i)' $\nu(A_i \setminus B) = 0$ for every i ;
- (ii) $\nu(B) = \lim_i \nu(A_i)$.

Property **AP**(null), which is a weak version of continuity from below, characterizes those finitely additive measures ν (defined on a σ -algebra \mathcal{F}), for which the space $L_1(\nu)$ is complete in the “usual” metric; see [5] and [4] (and see [1] for the theory of L_1 spaces over finitely additive measures).

The authors of [5] asked whether there exists a density with property **AP**(null). We give below a positive answer to that question and present some related results, some of which build on ideas from [7].

Every free ultrafilter \mathcal{U} on ω defines, in a natural way, a certain density $\nu_{\mathcal{U}}$ (see section 2). We shall show that $\nu_{\mathcal{U}}$ has property **AP**(null) whenever \mathcal{U} contains a set which is thin enough. Our next result yields a short proof of a result due to Mekler [7] — we show that $\nu_{\mathcal{U}}$ has property **AP**(*) provided \mathcal{U} is a P-point ultrafilter. We also prove that there is a \mathcal{U} giving a density $\nu_{\mathcal{U}}$ which has property **AP**(null) but fails to have **AP**(*). Finally, we explain that, roughly speaking, one cannot retrieve properties of \mathcal{U} from corresponding properties of $\nu_{\mathcal{U}}$. In particular, $\nu_{\mathcal{U}}$ can have property **AP**(*) even if \mathcal{U} is not a P-point.

2. Ultrafilters and densities.

Let \mathcal{U} be a free ultrafilter on a space X . Given a bounded function $\alpha : X \rightarrow \mathbf{R}$, we write $a = \mathcal{U}\text{-}\lim_x \alpha(x)$ if $\{x : |a_x - a| < \varepsilon\} \in \mathcal{U}$ for every positive ε . If \mathcal{U} is a free ultrafilter on ω then it is routine to check that the formula

$$\nu_{\mathcal{U}}(A) = \mathcal{U}\text{-}\lim_n \frac{|A \cap n|}{n},$$

defines a finitely additive extension of the asymptotic density to the power set of ω (that is, $\nu_{\mathcal{U}}$ is a density in our terminology). The averaging process involved in densities can also be expressed in terms of the Cesàro matrix, as explained in [7].

Recall that a free ultrafilter \mathcal{U} is called a *P-point* if for every sequence (X_i) of elements of \mathcal{U} one can find a set $Y \in \mathcal{U}$ which is almost contained in every X_i . It is well-known that P-points do exist under Martin’s axiom; however, their nonexistence is also relatively consistent, see [8, Section VI.4].

Given an infinite set $X \subseteq \omega$, we write

$$I_n^X = [\max(X \cap n), n) \cap \omega,$$

whenever $n \in X$. Say that a set X is *thin* if

$$\lim_{n \in X} \frac{|I_n^X|}{n} = 1.$$

In other words, a set X is thin if, enumerating X as $(n_k)_k$ in increasing order, we have $\lim_k n_k/n_{k+1} = 0$.

Lemma 1. *If an ultrafilter \mathcal{U} contains a thin set X then*

$$\nu_{\mathcal{U}}(A) = \mathcal{U}\text{-}\lim_n \frac{|A \cap I_n^X|}{n} = \mathcal{U}\text{-}\lim_n \frac{|A \cap I_n^X|}{|I_n^X|},$$

for every set A .

Proof. The second equation is obvious as X is thin and in \mathcal{U} . For the first, notice that

$$A \cap I_n^X \subseteq A \cap n \subseteq (A \cap I_n^X) \cup \max(X \cap n).$$

Therefore,

$$\frac{|A \cap I_n^X|}{n} \leq \frac{|A \cap n|}{n} \leq \frac{|A \cap I_n^X|}{n} + \frac{n - |I_n^X|}{n}.$$

Taking the limit along \mathcal{U} and remembering that $X \in \mathcal{U}$ is thin, we obtain the first equation of the lemma.

Theorem 1. *If an ultrafilter \mathcal{U} contains a thin set X then $\nu_{\mathcal{U}}$ is a density having property **AP**(null).*

Proof. Having an ultrafilter \mathcal{U} and a thin set $X \in \mathcal{U}$ fixed, we write $\nu = \nu_{\mathcal{U}}$ and $I_n = I_n^X$ for simplicity. By Lemma 1 we have $\nu(A) = \mathcal{U}\text{-}\lim_n \frac{|A \cap I_n|}{n}$ for every A .

To check **AP**(null) take an increasing sequence $(A_i)_{i \geq 1}$ and put $\alpha = \lim_i \nu(A_i)$. Passing to a subsequence we may assume that $\nu(A_i) \geq \alpha - 1/i$.

Find a decreasing sequence $(X_i)_{i \geq 1}$ of elements of \mathcal{U} such that $X_1 \subseteq X$, $X_i \cap i = \emptyset$, and

$$\left| \frac{|A_i \cap I_n|}{|I_n|} - \nu(A_i) \right| < \frac{1}{i},$$

whenever $n \in X_i$.

Now we define a set B , separately on each segment I_n . Put $B \cap I_n = \emptyset$ for $n \in X \setminus X_1$ and $B \cap I_n = A_i \cap I_n$ for $n \in X_i \setminus X_{i+1}$.

Let $n \in X_i$; then $n \in X_j \setminus X_{j+1}$ for some $j \geq i$. It follows that

$$\alpha - \frac{2}{i} \leq \nu(A_j) - \frac{1}{j} \leq \frac{|B \cap I_n|}{|I_n|} = \frac{|A_j \cap I_n|}{|I_n|} \leq \nu(A_j) + \frac{1}{j} \leq \alpha + \frac{1}{i},$$

and hence for every $n \in X_i$ we have

$$\left| \frac{|B \cap I_n|}{|I_n|} - \alpha \right| < \frac{2}{i},$$

which gives $\nu(B) = \alpha$.

Now we check that $\nu(A_i \setminus B) = 0$. Indeed, if $n \in X_i$ then for some $j \geq i$ we have $B \cap I_n = A_j \cap I_n \supseteq A_i \cap I_n$. Thus $(A_i \setminus B) \cap I_n = \emptyset$ for every $n \in X_i$, and we are done.

We next give a short proof of Mekler's result [7] that a P-point yields a density with **AP**(*).

Theorem 2. *If \mathcal{U} is a P-point ultrafilter then $\nu_{\mathcal{U}}$ is a density having property **AP**(*).*

Proof. For an increasing sequence $(A_i)_i$ we take sets X_i belonging to \mathcal{U} and such that

$$\left| \frac{|A_i \cap n|}{n} - \nu(A_i) \right| < \frac{1}{i},$$

whenever $n \in X_i$. As in the proof of Theorem 1, we write $\alpha = \lim_i \nu(A_i)$. Now, since \mathcal{U} is a P-point, there is a set $Y \in \mathcal{U}$ such that $Y \subseteq^* X_i$ for every i . Choose an increasing sequence of numbers $m_i \in \omega$ so that $Y \subseteq X_i \cup m_i$. We define a set B so that $B \cap [m_i, m_{i+1}) = A_i \cap [m_i, m_{i+1})$ for every i . We have $A_i \subseteq^* B$, since $A_i \setminus B \subseteq m_i$. If $n \in [m_i, m_{i+1})$ then $B \cap n \subseteq A_i \cap n$. If, moreover, $n \in Y$ then $n \in X_i$ so

$$\frac{|B \cap n|}{n} \leq \frac{|A_i \cap n|}{n} < \nu(A_i) + \frac{1}{i} \leq \alpha + \frac{1}{i}.$$

It follows that the inequality above holds for every $n \in Y \setminus m_i$, and therefore $\nu_{\mathcal{U}}(B) \leq \alpha$. On the other hand, the reverse inequality holds because $\nu_{\mathcal{U}}(B) \geq \nu_{\mathcal{U}}(A_i)$ for every i .

Theorem 3. *Suppose that \mathcal{U} is an ultrafilter containing a thin set. Then $\nu_{\mathcal{U}}$ has property **AP**(*) if and only if \mathcal{U} is a P-point.*

Proof. One direction is Theorem 2 (and doesn't need thinness). To prove the other direction, fix a thin set X belonging to \mathcal{U} ; again let I_n stand for I_n^X . Let $(X_m)_m$ be a decreasing sequence in \mathcal{U} . For every m write

$$A_m = \bigcup_{k \in X \setminus X_m} I_k.$$

Then $A_m \cap I_n = \emptyset$ for every $n \in X_m$, and Lemma 1 gives $\nu_{\mathcal{U}}(A_m) = 0$. Since $\nu_{\mathcal{U}}$ has property **AP**(*), there is a set A with $\nu_{\mathcal{U}}(A) = 0$, and such that $A_m \subseteq^* A$ for every m . It follows that

$$Y = \left\{ k : \frac{|A \cap I_k|}{|I_k|} < \frac{1}{2} \right\} \in \mathcal{U}.$$

Since A_m is almost contained in A , we have $A_m \subseteq A \cup k_m$ for some $k_m \in X$. Then $Y \subseteq X_m \cup k_m$. Indeed, if $k \in Y \setminus X_m$ then

$$\frac{|A \cap I_k|}{|I_k|} < \frac{1}{2} \text{ and } A_m \cap I_k = I_k.$$

Hence $A_m \cap I_k \not\subseteq A \cap I_k$, which means $k \leq k_m$. The proof is complete.

In connection with Theorem 3 it is perhaps worth remarking that, assuming Martin's axiom, one can easily construct a P-point which does not contain any thin set.

Given an arbitrary (infinite) thin set X , it is easy to find a free ultrafilter \mathcal{U} which contains X but is *not* a P-point (take a sequence of distinct ultrafilters containing X and let \mathcal{U} be its cluster point). In view of Theorem 2 and Theorem 3, this remark yields the following.

Corollary 4. *There exists a free ultrafilter \mathcal{U} such that $\nu_{\mathcal{U}}$ has property **AP**(null) but not property **AP**($*$).*

3. Different ultrafilters giving the same density.

Extending an idea suggested by Example 1.5 of [7], we shall show that rather dissimilar ultrafilters \mathcal{U} can lead to similar densities or even the same density.

If \mathcal{U} is an ultrafilter on a set X and $g : X \rightarrow Y$ is any mapping then $g(\mathcal{U})$ denotes the ultrafilter on Y consisting of those $B \subseteq Y$ for which $g^{-1}(B) \in \mathcal{U}$. Note that

$$g(\mathcal{U})\text{-}\lim_y \alpha(y) = \mathcal{U}\text{-}\lim_x \alpha(g(x)),$$

for every function α from Y into $[0, 1]$.

We consider the set $\Delta = \{(n, k) : n < k\}$ and "the standard pairing function" $p : \Delta \rightarrow \omega$, where

$$p(n, k) = \frac{k(k-1)}{2} + n.$$

Put also $q(k) = p(0, k)$ and denote by $\pi : \Delta \rightarrow \omega$ the projection onto the second coordinate.

Lemma 2. *If \mathcal{U} is a free ultrafilter on Δ then $\nu_{p(\mathcal{U})} = \nu_{q \circ \pi(\mathcal{U})}$.*

Proof. (1) Note first that for any $A \subseteq \omega$ and natural numbers Q, P , if $Q \leq P \leq Q + 2\sqrt{Q}$ then

$$\left| \frac{|A \cap Q|}{Q} - \frac{|A \cap P|}{P} \right| \leq \frac{2}{\sqrt{Q}}.$$

Indeed, (1) follows immediately from the following two inequalities:

$$0 \geq \frac{|A \cap Q|}{Q} - \frac{|A \cap P|}{Q} = -\frac{|A \cap (P \setminus Q)|}{Q} \geq -\frac{P - Q}{Q} \geq -\frac{2}{\sqrt{Q}};$$

$$0 \leq \frac{|A \cap P|}{Q} - \frac{|A \cap P|}{P} = |A \cap P| \left(\frac{1}{Q} - \frac{1}{P} \right) = \frac{|A \cap P|}{P} \cdot \frac{P - Q}{Q} \leq \frac{2}{\sqrt{Q}}.$$

(2) For every $n < k$ we have

$$q(k) \leq p(n, k) \leq q(k) + 2\sqrt{q(k)}.$$

Indeed, the definitions of $p(n, k)$ and $q(k)$ make the first inequality trivial, and they imply that

$$p(n, k) - q(k) = n \leq k - 1 \leq \sqrt{2q(k)} \leq 2\sqrt{q(k)},$$

which gives the second inequality.

(3) Using (2) and (1) (where $Q = q(k)$, $P = p(n, k)$), we get for any $A \subseteq \omega$

$$\begin{aligned} \nu_{p(\mathcal{U})}(A) &= p(\mathcal{U})\text{-}\lim_m \frac{|A \cap m|}{m} = \mathcal{U}\text{-}\lim_{(n,k)} \frac{|A \cap p(n, k)|}{p(n, k)} = \\ &= \mathcal{U}\text{-}\lim_{(n,k)} \frac{|A \cap q(k)|}{q(k)} = q \circ \pi(\mathcal{U}) \lim \frac{|A \cap m|}{m} = \nu_{q \circ \pi(\mathcal{U})}(A). \end{aligned}$$

Theorem 4. (a) *If there exist P-point ultrafilters on ω then there is an ultrafilter \mathcal{V} which is not a P-point, and such that $\nu_{\mathcal{V}}$ has property \mathbf{AP}^* .*
(b) *There is an ultrafilter \mathcal{V} which does not contain a thin set, and such that $\nu_{\mathcal{V}}$ has property $\mathbf{AP}(\text{null})$ but not \mathbf{AP}^* .*

Proof. Let \mathcal{U}_1 and \mathcal{U}_2 be ultrafilters on ω and let $\mathcal{U} = \mathcal{U}_1 \otimes \mathcal{U}_2$ denote their product. By definition, $D \in \mathcal{U}_1 \otimes \mathcal{U}_2$ if $\{n : D|_n \in \mathcal{U}_2\} \in \mathcal{U}_1$, where $D|_n$ is defined to be $\{k : (n, k) \in D\}$. Note that $\Delta \in \mathcal{U}_1 \otimes \mathcal{U}_2$, so we may consider the product ultrafilter as defined on Δ . Note also that $\pi(\mathcal{U}_1 \otimes \mathcal{U}_2) = \mathcal{U}_2$. To check (a) take a P-point \mathcal{U}_2 and arbitrary \mathcal{U}_1 , and consider $\mathcal{U} = \mathcal{U}_1 \otimes \mathcal{U}_2$. Using Lemma 2, we get

$$\nu_{p(\mathcal{U}_1 \otimes \mathcal{U}_2)} = \nu_{q \circ \pi(\mathcal{U}_1 \otimes \mathcal{U}_2)} = \nu_{q(\mathcal{U}_2)}.$$

Since $q(\mathcal{U}_2)$ is a P-point, $\nu_{p(\mathcal{U})}$ has property \mathbf{AP}^* . On the other hand, \mathcal{U} and its isomorphic copy $p(\mathcal{U})$ are not P-points, for no element of \mathcal{U} is almost contained in every $\Delta_N = \{(n, k) \in \Delta : n \geq N\}$.

We argue for (b) in a similar manner. Take an ultrafilter \mathcal{U}_2 which is not a P-point and contains a thin set. Then $q(\mathcal{U}_2)$ contains a thin set, so $\nu_{q(\mathcal{U}_2)}$ has property $\mathbf{AP}(\text{null})$. On the other hand, $q(\mathcal{U}_2)$ is not a P-point, so $\nu_{q(\mathcal{U}_2)}$ does not have property \mathbf{AP}^* by Theorem 3. Now $p(\mathcal{U})$ defines the same density but it does not contain a thin set. Indeed, if $A \subseteq \omega$ and $p^{-1}(A) \in \mathcal{U}$ then there are $i, j \in A$ with $i < j < 2i$.

4. Densities without additivity properties.

Recall that it is relatively consistent that no density has property \mathbf{AP}^* ; see Mekler [7], where Shelah's argument from [8] is suitably adapted. Frankiewicz, Shelah, Zbierski [3] obtained a model of set theory, in which there are no *ccc* P-sets in $\beta\omega \setminus \omega$. Let us note that this result improves Mekler's theorem. Indeed, every density ν defines the unique Radon measure $\hat{\nu}$ on the compact space $\beta\omega \setminus \omega$. If S denotes the support of $\hat{\nu}$ then S is clearly *ccc*. If, moreover, ν has property \mathbf{AP}^* then S is easily seen to be a P-set. Corollary 4 above shows that no extra axioms are needed to find an ultrafilter \mathcal{U} such that $\nu_{\mathcal{U}}$ fails to have \mathbf{AP}^* . We sketch here another, more constructive, argument for this fact.

(1) First note that there is a sequence (A_i) of subsets of ω such that for every k and $\varepsilon > 0$ the system of inequalities

$$\frac{|A_i \cap n|}{n} < \varepsilon, \quad i = 1, 2, \dots, k \quad \frac{|A_{k+1} \cap n|}{n} \geq \frac{1}{2},$$

is satisfied for infinitely many n .

This may be proved by a Baire category argument but, as the referee remarked, it suffices to put $A_i = \{[n_k, n_{k+1}) : k \in W_i\}$, where $(n_k)_k$ is an enumeration of a thin set, and $(W_i)_i$ is a partition of ω into infinite sets.

(2) Denote by \mathcal{B} the family of all sets B almost including A_i for every i . Put

$$X_{\varepsilon,i} = \left\{ n : \frac{|A_i \cap n|}{n} < \varepsilon \right\}; \quad Y_B = \left\{ n : \frac{|B \cap n|}{n} \geq \frac{1}{2} \right\},$$

for every i and $\varepsilon > 0$, and for every $B \in \mathcal{B}$.

(3) Note that the set

$$X_{\varepsilon,1} \cap X_{\varepsilon,2} \cap \dots \cap X_{\varepsilon,k} \cap Y_B$$

is infinite for any ε , k , and every $B \in \mathcal{B}$. Since $Y_B \cap Y_C \supseteq Y_{B \cap C}$ for $B, C \in \mathcal{B}$, it follows that the family of all $X_{\varepsilon,i}$ and all Y_B has the strong finite intersection property, and therefore is contained in some free ultrafilter \mathcal{U} . Now we have $\nu_{\mathcal{U}}(A_i) = 0$ for every i , and $\nu_{\mathcal{U}}(B) \geq 1/2$ for all $B \in \mathcal{B}$, so $\nu_{\mathcal{U}}$ does not have property \mathbf{AP}^* .

Of course, property $\mathbf{AP}(\text{null})$ is much harder to destroy, and we do not know if this can be done by a suitable modification of the argument above. In response to our question whether there is an ultrafilter giving a density without property $\mathbf{AP}(\text{null})$, David Fremlin presented the following result (in a letter of October, 1999).

Theorem 5 (Fremlin). *Suppose that \mathcal{U} is an ultrafilter on ω such that for every $A \in \mathcal{U}$ there is a $k > 0$ with $A + k \in \mathcal{U}$. Write $\mathcal{V} = g(\mathcal{U})$, where $g : \omega \rightarrow \omega$, $g(n) = 2^n$. Then the density $\nu_{\mathcal{V}}$ does not have property $\mathbf{AP}(\text{null})$.*

Proof. Clearly $\nu_\nu = \nu$, where ν is a density defined by the formula

$$\nu(A) = \mathcal{U}\text{-}\lim_n \frac{|A \cap 2^n|}{2^n}.$$

For every $k \in \omega$ we find a set $I_k \in \mathcal{U}$ such that $I_k \cap (k+1) = \emptyset$ and $|i-j| > k$ whenever $i, j \in I_k$, $i \neq j$. For every k we put

$$A_k = \bigcup_{i \in I_k} [2^{i-k-1}, 2^{i-k}) \cap \omega.$$

Note that $\nu(A_k) \leq 2^{-k}$. Indeed, if $i \in I_k$ then $A_k \cap [2^{i-k}, 2^i) = \emptyset$, so $|A_k \cap 2^i|/2^i \leq 2^{-k}$.

Now we check that, given a set B with $\nu(B) < 1/4$, we have $\nu(A_k \setminus B) > 0$ for infinitely many k . Let

$$J = \{i : |B \cap [2^{i-1}, 2^i)| \leq 2^{i-2}\}.$$

Then $J \in \mathcal{U}$, since otherwise we would have $\nu(B) \geq 1/4$.

It follows from the assumption on \mathcal{U} that $J+k \in \mathcal{U}$ for infinitely many k . Fix a number k with this property, put $K = (J+k) \cap I_k$, and consider any $i \in K$. Since $i \in I_k$ and $i-k \in J$, we have

$$A_k \supseteq [2^{i-k-1}, 2^{i-k}) \cap \omega, \quad |B \cap [2^{i-k-1}, 2^{i-k})| \leq 2^{i-k-2}.$$

Hence

$$|(A_k \setminus B) \cap 2^i| \geq |(A_k \setminus B) \cap [2^{i-k-1}, 2^{i-k})| \geq 2^{i-k} - 2^{i-k-1} - 2^{i-k-2} = 2^{i-k-2};$$

$$\frac{|(A_k \setminus B) \cap 2^i|}{2^i} \geq 2^{-(k+2)},$$

whenever $i \in K \in \mathcal{U}$. Hence $\nu(A_k \setminus B) > 0$. Therefore the sequence $A_4, A_4 \cup A_5, \dots$ witnesses that ν fails to have **AP**(null).

In order to get a density without property **AP**(null), it is now sufficient to find an ultrafilter \mathcal{U} such that for every $A \in \mathcal{U}$ there is a $k > 0$ with $A+k \in \mathcal{U}$. D. Fremlin pointed out that the existence of an ultrafilter with this property may be quickly derived from Glazer's theorem on idempotent ultrafilters (see e.g. [9], section 15). This suggested us the following straightforward argument.

The formula $\phi(\mathcal{X}) = \{A : A+1 \in \mathcal{X}\}$ defines a continuous mapping $\phi : \beta\omega \setminus \omega \rightarrow \beta\omega \setminus \omega$. By compactness, there is a minimal ϕ -invariant closed and nonempty subset S of $\beta\omega \setminus \omega$. We check that any $\mathcal{U} \in S$ has the required property. Indeed, the orbit $\{\phi^k(\mathcal{U}) : k \geq 1\}$ must be dense in S , since its closure is ϕ -invariant. Therefore, given $A \in \mathcal{U}$, there is a $k \geq 1$ such that the ultrafilter $\phi^k(\mathcal{U})$ is in the closed and open set defined by A . Hence $A \in \phi^k(\mathcal{U})$, so $A+k \in \mathcal{U}$, and we are done.

Acknowledgements. We would like to thank the referee for a careful reading of the first version of this paper, and for several valuable comments. Thanks are also due to David Fremlin for sending us his result, and for his kind permission to include it here.

References

- [1] K.P.S. Bhaskara Rao, M. Bhaskara Rao, *Theory of charges*, Academic Press (1983).
- [2] R.C. Buck, *Generalized asymptotic density*, Amer. J. Math. 75 (1953), 335–346.
- [3] R. Frankiewicz, S. Shelah, P. Zbierski, *On closed P -sets with ccc in the space ω^** , J. Symbolic Logic 58 (1993), 1171–1176.
- [4] S. Gangopadhyay, *On the completeness of L_p spaces over a charge*, Colloq. Math. 58 (1990), 291–300.
- [5] S. Gangopadhyay, B.V. Rao, *Completeness of L_p spaces over finitely additive measures*, Colloq. Math. 80 (1999), 83–95.
- [6] D. Maharam, *Finitely additive measures on the integers*, Sankhya Ser. A 38 (1976), 44–59.
- [7] A.H. Mekler, *Finitely additive measures on N and the additive property*, Proc. Amer. Math. Soc. 92 (1984), 439–444.
- [8] S. Shelah, *Proper forcing*, Lecture Notes in Mathematics 940, Springer-Verlag, Berlin 1982.
- [9] S. Todorcevic, *Topics in topology*, Springer (1997), Berlin.

Department of Mathematics, University of Michigan

e-mail: ABLASS@MATH.LSA.UMICH.EDU

Institute of Mathematics, Polish Academy of Sciences

e-mail: RF@IMPAN.GOV.PL

Institute of Mathematics, University of Wrocław, pl. Grunwaldzki 2/4, 50–218 Wrocław, Poland

e-mail: GRZES@MATH.UNI.WROC.PL

*Institute of Mathematics, Wrocław Technical University
and Institute of Mathematics, Polish Academy of Sciences*

e-mail: CRN@GRAF.IM.PWR.WROC.PL