

INACCESSIBLE CARDINALS WITHOUT THE AXIOM OF CHOICE

ANDREAS BLASS, IOANNA M. DIMITRIOU, BENEDIKT LÖWE

ABSTRACT. We consider four notions of strong inaccessibility that are equivalent in ZFC and show that they are not equivalent in ZF.

1. INTRODUCTION

Large cardinals serve as a calibration of logical strength for axiom systems of set theory. If we compare theories up to equiconsistency, every natural set-theoretic axiom system is equiconsistent with some $\text{ZFC} + \text{L}$ where L is a large cardinal axiom. These axioms come in many guises, and most of the large cardinal axioms have several equivalent definitions. For instance, a cardinal κ is measurable if there is a κ -complete nonprincipal ultrafilter on it, but in the presence of the axiom of choice you can show that this is equivalent to the statement “there is an elementary embedding of the universe into a transitive class with critical point κ ”.

If you move into the choiceless world, some of these ZFC-equivalent definitions are not equivalent anymore. The cardinal \aleph_1 can never be the critical point of an elementary embedding of the universe into a transitive class, but it is possible that it carries a σ -complete nonprincipal ultrafilter in models of ZF.¹ It is interesting to analyze how much of the robustness of large cardinal notions we lose if we give up the axiom of choice.

In this paper, we investigate a large cardinal notion that is in conflict with negations of the axiom of choice: the notion of strong inaccessibility. A strongly inaccessible cardinal κ is a regular strong limit cardinal where the traditional ZFC-definition of “strong limit” is that for all $\lambda < \kappa$, we have $2^\lambda < \kappa$.

2000 *Mathematics Subject Classification.* **03E10 03E25** 03E55.

¹*Cf.* [Je03, Theorem 21.16]. Even worse, in some cases, even the consistency strength of $\text{ZFC} + \text{L}$ and $\text{ZF} + \text{L}$ is different. For instance, let L be “there is a cardinal κ such that $\mathcal{P}_\kappa(\kappa^+)$ carries a normal measure”. Then the consistency strength of $\text{ZF} + \text{L}$ is bounded from above by ω Woodin cardinals ([Be81] in combination with Woodin’s result on the consistency strength of AD; *cf.* [Ka03, Theorem 32.16]) whereas the consistency strength of $\text{ZFC} + \text{L}$ is far bigger. (The last statement combines a result of Solovay on the failure of \square_κ at a κ^+ -supercompact cardinal κ with inner model results by Schimmerling and Zeman [ScZe01]. For more details, *cf.* [Bo02, Kapitel 6].)

In the ZF-context, this doesn't necessarily make sense, as 2^λ may not be well-ordered. We look at four different definitions of inaccessible cardinals that are equivalent in ZFC but not equivalent in ZF (*i*-inaccessibility, *v*-inaccessibility, \bar{s} -inaccessibility, and \bar{v} -inaccessibility) and study the implicational structure of these notions (in ZF).

The paper is organized as follows: In §2, we present the definitions and their simple consequences. The four mentioned notions are presented in an implication diagram in Figure 1. In §3, we first finish the proof of the positive part of the implication diagram. After that, we briefly digress and discuss a property $(*\omega_1, \bar{v})$ that corresponds to being inaccessible without being a limit cardinal in §4. The main part of the paper contains the three independence results proving that none of the arrows in Figure 1 are reversible in §§5, 6, and 7. Throughout this paper, the ambient theory is ZF.

2. DEFINITIONS OF STRONG LIMITS

Ordinarily, in set theory with the axiom of choice, we write $X \leq Y$ to mean that there is an injection from X to Y , or equivalently (provided $X \neq \emptyset$) that there is a surjection from Y to X . Without the axiom of choice, these two statements are not equivalent anymore and we have to be much more precise. We define four different notions of ‘being smaller than’ for the context of ZF:

Let X and Y be nonempty sets and α be an ordinal. Then we write

- $X <_i \alpha$ if there is some $\beta < \alpha$ and an injection from X into β ,
- $X <_s \alpha$ if there is some $\beta < \alpha$ and a surjection from β onto X ,
- $X <_{\bar{v}} Y$ if there is no injection from Y into X , and
- $X <_{\bar{s}} Y$ if there is no surjection from X onto Y .

Of course, if X and Y are wellorderable, these four relations coincide. The following are immediate observations:

Observation 1. For a set X and an ordinal α , the statements $X <_s \alpha$ and $X <_i \alpha$ are equivalent and both imply that X is wellorderable.

Observation 2. If κ is a cardinal, then $X <_s \kappa$ implies $X <_{\bar{s}} \kappa$.

Observation 3. For sets X and Y , if $X <_{\bar{s}} Y$, then $X <_{\bar{v}} Y$.

In the following, we shall use the symbol \mathbf{x} as a placeholder for one of the four symbols *i*, *s*, \bar{v} and \bar{s} . We say that a limit cardinal κ is an **x-strong limit** if for all $\lambda < \kappa$, we have $\mathcal{P}(\lambda) <_{\mathbf{x}} \kappa$, and we call an uncountable cardinal κ **x-inaccessible** if it is a regular **x-strong limit** cardinal.²

Of course, inaccessibility of a cardinal κ is typically connected with metamathematical properties of \mathbf{V}_κ . This suggests another natural definition of inaccessibility: An ordinal κ is **v-inaccessible** if \mathbf{V}_κ is a model of second-order ZF. Here, by \mathbf{V}_κ we mean the collection of sets of (Mirimanoff) rank

²Here and for the rest of the paper, ‘cardinal’ means ‘initial ordinal’.




FIGURE 1. The implication diagram

$< \kappa$, regarded as a structure by interpreting \in as actual membership. By “second-order ZF”, we mean ZF with replacement strengthened to say that the image of a set under any function (not just under definable functions) is a set. It is easy to check the following characterizations of v -inaccessibility.

Proposition 4. The following are equivalent, for any ordinal κ .

- (1) κ is v -inaccessible,
- (2) κ is a regular, uncountable cardinal and for all $\alpha < \kappa$, we have $\mathbf{V}_\alpha <_{\bar{s}} \kappa$, and
- (3) κ is a regular, uncountable cardinal and for all $\alpha < \kappa$ and all cofinal subsets $C \subseteq \kappa$, we have $\mathbf{V}_\alpha <_{\bar{s}} C$.

Combining Proposition 4 with the fact that $\mathcal{P}(\lambda) \subseteq \mathbf{V}_{\lambda+1}$, we immediately get the following connection between v -inaccessibility and \bar{s} -inaccessibility:

Corollary 5. Every v -inaccessible cardinal is \bar{s} -inaccessible.

The implication diagram in Figure 1 gives the connections between the five notions of inaccessibility. All implications except for the ones labelled (‡) follow from the simple observations and Corollary 5. The implication (‡) will be proved as Theorem 11.

It is quite easy to see that the notion of an i -strong limit is too strong for purposes of set theory without the axiom of choice:

Proposition 6. If there is an uncountable i -strong limit, then there is a wellordering of the real numbers.

Proof. If κ is uncountable, then $\aleph_0 < \kappa$, so by Observation 1, $\mathcal{P}(\omega)$ is wellorderable. q.e.d.

We now use the notation IC for the assertion “there is an inaccessible cardinal” (meaning “there is an i -inaccessible cardinal” if we use it in the $\neg\text{AC}$ -context). For the separating instances (without AC), we use the following notation:

$\text{IC}_{\mathbf{v}/\neg\mathbf{i}}$ “there is an \mathbf{v} -inaccessible cardinal which is not \mathbf{i} -inaccessible”,
 $\text{IC}_{\bar{\mathbf{s}}/\neg\mathbf{v}}$ “there is an $\bar{\mathbf{s}}$ -inaccessible cardinal which is not \mathbf{v} -inaccessible”,
 $\text{IC}_{\bar{\mathbf{i}}/\neg\bar{\mathbf{s}}}$ “there is an $\bar{\mathbf{i}}$ -inaccessible cardinal which is not $\bar{\mathbf{s}}$ -inaccessible”.

The goal of this paper is to show that the arrows in the above diagram do not reverse in \mathbf{ZF} , *i.e.*, that $\mathbf{ZF} + \text{IC}_{\mathbf{v}/\neg\mathbf{i}}$, $\mathbf{ZF} + \text{IC}_{\bar{\mathbf{s}}/\neg\mathbf{v}}$, and $\mathbf{ZF} + \text{IC}_{\bar{\mathbf{i}}/\neg\bar{\mathbf{s}}}$ are consistent (*cf.* Theorems 17, 20, and 25). On the metamathematical side, it is interesting to note that the different definitions of strong inaccessibility do not give different consistency strength:

Proposition 7. If κ is $\bar{\mathbf{i}}$ -inaccessible, then κ is inaccessible in \mathbf{L} .

Proof. Clearly, κ is regular in \mathbf{L} . Let $\alpha < \kappa$. If $\mathbf{L} \models$ “there is an injection from κ into $\mathcal{P}(\alpha)$ ”, then there is an injection from κ into $\mathcal{P}^{\mathbf{L}}(\alpha) \subseteq \mathcal{P}(\alpha)$, so κ is not $\bar{\mathbf{i}}$ -inaccessible in the universe. q.e.d.

Corollary 8. The following are equiconsistent:

- (1) $\text{ZFC} + \text{IC}$,
- (2) $\text{ZF} +$ “there is an \mathbf{i} -inaccessible”,
- (3) $\text{ZF} +$ “there is an \mathbf{v} -inaccessible”,
- (4) $\text{ZF} +$ “there is an $\bar{\mathbf{s}}$ -inaccessible”, and
- (5) $\text{ZF} +$ “there is an $\bar{\mathbf{i}}$ -inaccessible”.

This project of this paper was in part motivated by the behaviour of cardinals under the assumption of $\text{ZF} + \text{AD}$ where AD denotes the axiom of determinacy (*cf.* [Ka03, § 27]).³

In the ZFC -context, the property

$$(*_{\kappa, \mathbf{x}}) \quad \text{for all } \lambda < \kappa, \text{ we have } \mathcal{P}(\lambda) <_{\mathbf{x}} \kappa$$

can only hold for limit cardinals. This remains true without the axiom of choice for $<_{\bar{\mathbf{s}}}$:

Observation 9. The statement $(*_{\kappa^+, \bar{\mathbf{s}}})$ is inconsistent with ZF .

Proof. In ZF , there is a surjection from $\mathcal{P}(\kappa)$ onto κ^+ . q.e.d.

The analogue of Observation 9 for $\bar{\mathbf{i}}$ is not true: By a theorem of Mycielski’s, AD implies that there cannot be an injection from ω_1 into the reals [Ka03, Proposition 27.11 (a)]; therefore $(*_{\omega_1, \bar{\mathbf{i}}})$ holds under AD . We’ll return to $(*_{\omega_1, \bar{\mathbf{i}}})$ in § 4.

³For an alternative definition of strong inaccessibility designed specifically for the AD -context, *cf.* [Ke85]. There, a cardinal κ is strongly inaccessible if it is regular and for all $\lambda < \kappa$ there is a bijection between the set of ultrafilters on λ and a proper initial segment of κ . Kechris proved (under the assumption of $\text{AD} + \mathbf{V} = \mathbf{L}(\mathbb{R})$) that δ_1^2 is strongly inaccessible in this sense.

3. \mathbf{v} -INACCESSIBLE CARDINALS.

In this section, we shall show the remaining implication (‡) from our diagram of implications. We start with a simple fact:

Lemma 10. Let κ be \mathbf{i} -inaccessible. If $\alpha < \kappa$ and \mathbf{V}_α is well-orderable, then $\mathbf{V}_\alpha <_{\mathbf{i}} \kappa$.

Proof. Suppose not, and let α be the first counterexample. By \mathbf{i} -inaccessibility of κ , this α cannot be a successor ordinal, and it obviously isn't 0, so it is a limit ordinal. By minimality of α , we have $\mathbf{V}_\beta <_{\mathbf{i}} \kappa$ for all $\beta < \alpha$. By regularity of κ , find a cardinal $\lambda < \kappa$ such that for all $\beta < \alpha$, we have $\mathbf{V}_\beta <_{\mathbf{i}} \lambda$.

Use the hypothesis to fix a well-ordering of \mathbf{V}_α ; it restricts to well-orderings of \mathbf{V}_β for all $\beta < \alpha$, which then determine order-isomorphisms $f_\beta : \mathbf{V}_\beta \rightarrow \tau_\beta$, where, by our choice of λ , we have $\tau_\beta < \lambda$ for all $\beta < \alpha$. Combining all these maps f_β , we get a one-to-one map $g : \mathbf{V}_\alpha \rightarrow \alpha \times \lambda$ defined by

$$g(x) := \langle \varrho(x), f_{\varrho(x)}(x) \rangle,$$

where $\varrho(x)$ is the Mirimanoff rank of x . Since $\alpha \times \lambda$ is well-orderable with order-type $< \kappa$, this finishes the proof. q.e.d.

Theorem 11. Every \mathbf{i} -inaccessible cardinal is \mathbf{v} -inaccessible.

Proof. Let κ be \mathbf{i} -inaccessible. By definition, κ is regular and uncountable, so it remains to prove that no \mathbf{V}_α with $\alpha < \kappa$ can be mapped onto κ . In fact, we shall prove more, namely that every such \mathbf{V}_α is well-orderable and has cardinality $< \kappa$.

By Lemma 10, it remains to show that \mathbf{V}_α is well-orderable for all $\alpha < \kappa$. Suppose not, and let α be the smallest counterexample. So \mathbf{V}_β is well-orderable and, by Lemma 10, of cardinality $< \kappa$ for all $\beta < \alpha$. As in the proof of Lemma 10, α must be a limit ordinal, and we can obtain a cardinal $\lambda < \kappa$ such that $\mathbf{V}_\beta <_{\mathbf{i}} \lambda$ for all $\beta < \alpha$. Thus, each such \mathbf{V}_β has a well-ordering of length $< \lambda$.⁴

By \mathbf{i} -inaccessibility of κ , fix a well-ordering \prec of $\mathcal{P}(\lambda)$. We inductively define, for $\beta \leq \alpha$, specific well-orderings \prec_β of \mathbf{V}_β as follows. Note that the induction includes $\beta = \alpha$, so we will obtain a well-ordering \prec_α of \mathbf{V}_α , thereby completing the proof.

Let \prec_0 be the empty relation (on the empty set $\mathbf{V}_0 = \emptyset$). For limit β , let

$$x \prec_\beta y : \iff \varrho(x) < \varrho(y) \text{ or } (\varrho(x) = \varrho(y) = \gamma \text{ and } x \prec_{\gamma+1} y).$$

Finally, for a successor ordinal $\beta = \gamma + 1$, notice that the order-isomorphism from \prec_γ to its order-type τ maps \mathbf{V}_γ one-to-one into λ (as our choice of λ ensures $\tau < \lambda$). This induces a one-to-one map of $\mathbf{V}_\beta = \mathcal{P}(\mathbf{V}_\gamma)$ into $\mathcal{P}(\lambda)$.

⁴If we could *choose* one such well-ordering for each $\beta < \alpha$, then we could combine these, as in the proof of the lemma, to well-order \mathbf{V}_α . Unfortunately, we cannot immediately make such choices. The rest of the proof is devoted to getting such choices without invoking the axiom of choice.

Use this map to pull back the fixed well-ordering of $\mathcal{P}(\lambda)$ to a well-ordering of \mathbf{V}_β , and define that well-ordering to be \prec_β . q.e.d.

4. THE PROPERTY $(*\omega_1, \bar{\delta})$

In this section, we briefly return to the property $(*\kappa, \mathbf{x})$ for successor cardinals κ . We'll discuss the case $\kappa = \omega_1$. We have seen that $(*\omega_1, \bar{\delta})$ is refutable in \mathbf{ZF} , but $(*\omega_1, \bar{\delta})$ is a theorem of $\mathbf{ZF} + \mathbf{AD}$. Even more, $\mathbf{ZF} + (*\omega_1, \bar{\delta})$ is equiconsistent with \mathbf{ZF} . For this, we give the following simple argument using the Feferman-Lévy model which is a folklore result well-known in the community:

The Feferman-Lévy model (*cf.* [FeLé63] and [Je03, Example 15.57]) is a symmetric submodel of a generic extension that satisfies “ \mathbb{R} is a countable union of countable sets” (–38 in the notation of [HoRu98]).

Lemma 12. If X is a countable union of countable sets, then there is no surjection from X onto $\aleph_1^{\aleph_0}$.

Proof. This is essentially the diagonal argument for König's Theorem: Let $X = \bigcup_{n \in \omega} X_n$ for some countable sets X_n and let π be any function from X to $\aleph_1^{\aleph_0}$. The set $\pi[X_n]$ is a countable set, and therefore $\omega_1 \setminus \{\pi(x)(n); x \in X_n\} \neq \emptyset$. Let α_n be the least element of this set. Define $Z : \omega \rightarrow \omega_1$ by $Z(n) := \alpha_n$. Then Z is not in the range of π . q.e.d.

Lemma 13. If there is an injection from ω_1 into the reals (170 in the notation of [HoRu98]), then there is a bijection between the reals and $\aleph_1^{\aleph_0}$.

Proof. By the assumption, we have an injection from \aleph_1 into 2^{\aleph_0} , and hence from $\aleph_1^{\aleph_0}$ into $(2^{\aleph_0})^{\aleph_0}$ and thus into 2^{\aleph_0} . By the Schröder-Bernstein theorem, we get a bijection from 2^{\aleph_0} to $\aleph_1^{\aleph_0}$. q.e.d.

Obviously, Lemmas 12 and 13 show that the Feferman-Lévy model satisfies $(*\omega_1, \bar{\delta})$.

Corollary 14. If $\text{Cons}(\mathbf{ZFC})$, then $\text{Cons}(\mathbf{ZF} + (*\omega_1, \bar{\delta}))$.

5. CARDINALS THAT ARE NOT $\dot{\mathbf{i}}$ -INACCESSIBLE.

For the remainder of the paper, we shall freely use the notation for forcing and symmetric submodels of generic extensions as laid out in [Je03, p. 249-261]. In the \mathbf{ZFC} context, we can recursively define for any cardinal μ its **beth sequence** by $\beth_0(\mu) := \mu$, $\beth_{\alpha+1}(\mu) := 2^{\beth_\alpha(\mu)}$, and $\beth_\delta(\mu) := \bigcup\{\beth_\alpha(\mu); \alpha < \delta\}$ (for limit ordinals δ).

Lemma 15. Let M be a model of \mathbf{ZFC} . Assume that, in M , λ , μ , and κ are cardinals satisfying $2^\lambda < \kappa$, $2^\mu < \kappa$. Assume further that, in M , \mathbb{P} is a partial order with $\text{Card}(\mathbb{P}) = \mu$. Let G be \mathbb{P} -generic over M , and $M \subseteq N \subseteq M[G]$ such that N is a model of \mathbf{ZF} . Then

$$N \models 2^\lambda <_{\bar{\delta}} \kappa.$$

Similarly, if in addition, in M , $\beth_\lambda(\mu) < \kappa$, then

$$N \models \mathbf{V}_\lambda <_{\bar{s}} \kappa.$$

Proof. This lemma is a standard and basic technique of forcing. Let us call a \mathbb{P} -name τ **canonical** if it consists of pairs $\langle \check{\alpha}, p \rangle$ for $\alpha < \lambda$. Obviously, there are at most $(2^{\text{Card}(\lambda \times \mathbb{P})})^M$ many canonical names. In $M[G]$, every subset of λ has a canonical name, so we have

$$(2^\lambda)^{M[G]} \leq (2^{\text{Card}(\lambda \times \mathbb{P})})^M = (2^{\max\{\lambda, \mu\}})^M < \kappa.$$

Therefore, $M[G] \models$ “there is no surjection from 2^λ onto κ ”.

Suppose towards a contradiction, N has a surjection from $(2^\lambda)^N$ onto κ . This persists in $M[G]$ and (since $(2^\lambda)^N \subseteq (2^\lambda)^{M[G]}$) can be trivially extended to a surjection from 2^λ onto κ in $M[G]$. Contradiction!

For the additional claim, we define **canonical α -names** for elements of \mathbf{V}_α by recursion: Elements F of \mathbf{V}_ω are absolute and can be represented by the name \check{F} . These names are the canonical 0-names. If C is the set of canonical α -names (for elements of $\mathbf{V}_{\omega+\alpha}$), then a canonical $\alpha+1$ -name is a subset of $C \times \mathbb{P}$. Recursively, it is easy to see that there are at most $\beth_\lambda(\mu)^M$ many canonical λ -names, so $M[G] \models$ “there is no surjection from \mathbf{V}_λ onto κ ” which again persists downwards to N . q.e.d.

Corollary 16. Let M be a model of ZFC+IC and let $\kappa \in M$ be inaccessible. If $\mathbb{P} \in M$ is a partial order with $M \models \text{Card}(\mathbb{P}) < \kappa$ and N is a ZF-model such that $M \subseteq N \subseteq M[G]$, then

$$N \models “\kappa \text{ is } \mathbf{v}\text{-inaccessible}”.$$

Proof. Since $\text{Card}(\mathbb{P}) < \kappa$, κ remains regular in $M[G]$ and hence in N . The rest follows from Lemma 15. q.e.d.

Theorem 17. If $\text{Cons}(\text{ZFC} + \text{IC})$, then $\text{Cons}(\text{ZF} + \text{IC}_{\mathbf{v}/\neg i})$.

Proof. Assume that κ is strongly inaccessible in $M \models \text{ZFC}$. If \mathbb{P} is Cohen forcing and G is \mathbb{P} -generic over M , we can create a symmetric submodel $N \subseteq M[G]$ such that the reals in N are not wellordered (*cf.* [Je03, Example 15.52]). By Proposition 6, we know that $N \models “\kappa$ is not \mathbf{i} -inaccessible”. But by Corollary 16, $N \models “\kappa$ is \mathbf{v} -inaccessible”. q.e.d.

6. CARDINALS THAT ARE NOT \mathbf{v} -INACCESSIBLE.

In this section, we'll construct a model of $\text{ZF} + \text{IC}_{\bar{s}/\neg \mathbf{v}}$. We start with a ground model M satisfying $\text{ZFC} + “\text{there exists an inaccessible}”$ and let κ be a fixed inaccessible cardinal in M . Consider the partial order

$$\mathbb{P} := \text{Fn}(\kappa \times \omega, 2)$$

adding a κ -sequence of Cohen reals. Let \mathbf{G} be the group of all permutations of κ , acting on \mathbb{P} by

$$\pi(p)(\alpha, n) := p(\pi^{-1}(\alpha), n);$$

it permutes the canonical names \dot{c}_α of the Cohen reals by acting on the subscripts α . If $A \subseteq \kappa$ is a set, then let $K_A \subseteq \mathbf{G}$ be the set of those permutations that fix the elements of A . Now take the filter \mathcal{F} on \mathbf{G} generated by the set $\{K_A; A \subseteq \kappa, \text{Card}(A) < \kappa\}$. Fix a \mathbb{P} -generic filter G over M and let N be the symmetric submodel of $M[G]$ defined by \mathcal{F} .

Clearly, $c_\alpha \in N$, as $K_{\{\alpha\}}$ is a support of its canonical name. For each $\lambda < \kappa$, the initial segment $C_\lambda := \langle c_\alpha; \alpha < \lambda \rangle$ of the adjoined κ -sequence of Cohen reals is in N , because its canonical name is supported by K_λ .

Proposition 18. In N , there is a surjection from $\mathbf{V}_{\omega+4}$ onto κ ; in particular, κ is not v -inaccessible (by Lemma 4).

Proof. Consider, in N , the function t assigning to every well-ordering (in N) of a subset (in N) of $\mathcal{P}^N(\omega)$ its order-type. For every $\lambda < \kappa$, the set $C_\lambda \in N$ witnesses that $\lambda \in \text{ran}(t)$. But the domain of t is a subset of $\mathbf{V}_{\omega+4}$. q.e.d.

Proposition 19. In N , κ is \bar{s} -inaccessible.

Proof. Consider $\lambda < \kappa$ and suppose, toward a contradiction, that N contains a map of $\mathcal{P}^N(\lambda)$ onto κ . Fix, in the ground model M , a name \dot{f} and a condition p such that $p \Vdash \text{“}\dot{f} : \mathcal{P}^N(\lambda) \rightarrow \kappa \text{ is surjective”}$.

For the rest of this proof, we shall work in M . Fix $\mu < \kappa$ large enough so that K_μ supports \dot{f} and so that $\text{dom}(p) \subseteq \mu \times \omega$.

Fix some $\alpha < \kappa$. Since p forces α to be in the range of \dot{f} , we can choose a nice name \dot{x}_α for a subset of λ and choose a condition q_α extending p and forcing that $\dot{f}(\dot{x}_\alpha) = \alpha$. Thanks to the countable chain condition of \mathbb{P} , there is a set $I \subseteq \kappa$ such that $\text{Card}(I) \leq \lambda$, $\text{dom}(q_\alpha) \subseteq I \times \omega$, and $\text{dom}(r) \subseteq I \times \omega$ for every condition r involved in the nice name \dot{x}_α . (Note that the conditions involved in a nice name for a subset of λ constitute λ antichains, each of which is countable.) Let $\pi_\alpha \in \mathbf{G}$ be a permutation that fixes all ordinals $< \mu$ but moves all ordinals in $I \setminus \mu$ to ordinals between μ and $\mu + \lambda$ (ordinal sum). Then π_α fixes \dot{f} and p , by our choice of μ . Therefore,

- (1) $\pi_\alpha(\dot{x}_\alpha)$ is a nice name for a subset of λ , and all conditions involved in it have domains $\subseteq (\mu + \lambda) \times \omega$.
- (2) $\pi_\alpha(q_\alpha)$ is a condition extending p and having domain $\subseteq (\mu + \lambda) \times \omega$.
- (3) $\pi_\alpha(q_\alpha) \Vdash \dot{f}(\pi_\alpha(\dot{x}_\alpha)) = \alpha$.

Now let α vary over all ordinals $< \kappa$. The number of possibilities for $\pi_\alpha(\dot{x}_\alpha)$ and $\pi_\alpha(q_\alpha)$ are, thanks to the $\mu + \lambda$ bounds in (1) and (2), strictly smaller than κ , because κ is inaccessible (in M). So there must be some $\alpha < \beta < \kappa$ with $\pi_\alpha(\dot{x}_\alpha) = \pi_\beta(\dot{x}_\beta)$ and $\pi_\alpha(q_\alpha) = \pi_\beta(q_\beta)$. But then (3) and the corresponding assertion for β say that one and the same condition forces one and the same term to have two distinct values, a contradiction. q.e.d.

Theorem 20. If $\text{Cons}(\text{ZFC} + \text{IC})$, then $\text{Cons}(\text{ZF} + \text{IC}_{\bar{s}/-v})$.

7. CARDINALS THAT ARE NOT \bar{s} -INACCESSIBLE.

In this last section, we'll construct a model of $\text{ZF} + \text{IC}_{\bar{\tau}/-\bar{s}}$. We again start with a ground model M satisfying $\text{ZFC} +$ "there exists an inaccessible" and let κ be a fixed inaccessible cardinal in M . Here we consider the partial order

$$\mathbb{P} := \text{Fn}(\kappa \times \omega \times \omega, 2)$$

which adds for each $\alpha < \kappa$ a countable set of Cohen reals $\{c(\alpha, n); n \in \omega\}$. We are going to permute these sets so that we cannot pick from these countable sets.

Let \mathbf{G} be the group of permutations of $\kappa \times \omega$ that fix the first component, *i.e.*, the direct product of κ copies of the group of all permutations of ω . Let this group act on \mathbb{P} by permuting the indices $\langle \alpha, n \rangle$ of the Cohen reals, *i.e.*,

$$\pi(p)(\alpha, n, m) := p(\pi^{-1}(\alpha, n), m).$$

Similarly to § 6, we let $K_A \subseteq \mathbf{G}$ be the set of those permutations that fix the elements of $A \subseteq \kappa \times \omega$. This time, we take the filter \mathcal{F} on \mathbf{G} generated by the set $\{K_A; A \subseteq \kappa \times \omega \text{ finite}\}$. Fix a \mathbb{P} -generic filter G over M and let N be the symmetric submodel of $M[G]$ defined by \mathcal{F} . Clearly, $c(\alpha, n) \in N$, as $K_{\{\langle \alpha, n \rangle\}}$ is a support of its canonical name.

Lemma 21. In N , the cardinal κ is not \bar{s} -inaccessible.

Proof. It is enough to show that there is a surjection from the reals onto κ in N . We consider the function $f \in M[G]$ defined by

$$f(x) := \begin{cases} \alpha & \text{if } x = c(\alpha, n) \text{ for some } n, \text{ or} \\ 0 & \text{otherwise.} \end{cases}$$

It is easy to see that f has a hereditarily symmetric name, and thus $f \in N$. q.e.d.

We now turn to the proof that κ is $\bar{\tau}$ -inaccessible in N . For $A \subseteq \kappa \times \omega$, $p \in \mathbb{P}$ and $J \subseteq \mathbb{P}$, we define

$$\begin{aligned} p^A &:= \{\langle \alpha, n, m, i \rangle \in p; \langle \alpha, n \rangle \in A\} = p \cap (A \times \omega \times 2), \text{ and} \\ J^A &:= \{p^A; p \in J\}. \end{aligned}$$

Lemma 22. Fix some finite $A \subseteq \kappa \times \omega$. Let φ be a formula of the forcing language such that for all names τ occurring in φ , we have that $K_A \subseteq \text{sym}(\tau)$. Then the following holds for all $p \in \mathbb{P}$:

$$\text{If } p \Vdash \varphi, \text{ then } p^A \Vdash \varphi.$$

Proof. If $p^A \not\Vdash \varphi$, then there is a $q \supseteq p^A$ such that $q \Vdash \neg\varphi$. Find a permutation $\pi \in K_A$ such that $\pi(p)$ and q are compatible, *i.e.*, there is an r such that $r \leq \pi(p)$ and $r \leq q$. Then $r \Vdash \neg\varphi$. Clearly, $\pi(p)$ forces φ^π , the formula obtained from φ by replacing all names occurring in φ by their π -images. Since $\pi \in K_A$, and by assumption all of the names in φ are supported by K_A , we get that $\varphi^\pi = \varphi$, so $r \Vdash \varphi$ which is absurd. q.e.d.

We shall now look at intermediate models between M and $M[G]$. For some finite $A \subseteq \kappa \times \omega$, we consider the forcing notion $\mathbb{P}^A = \{p^A; p \in \mathbb{P}\}$ which is forcing equivalent to ordinary Cohen forcing. Clearly, $G^A = \{p^A; p \in G\}$ is a \mathbb{P}^A -generic filter over M , and $M[G^A]$ is essentially the Cohen model.

Lemma 23. If $X \in N$ is a set of ordinals, then there is a finite $A \subseteq \kappa \times \omega$ such that $X \in M[G^A]$.

Proof. Since $X \in N$, it has a hereditarily symmetric \mathbb{P} -name \dot{X} (with respect to G). Fix a finite set $A \subseteq \kappa \times \omega$ such that $K_A \subseteq \text{sym}(\dot{X})$. Let $\varphi_\alpha \simeq \check{\alpha} \in \dot{X}$. Clearly, for each α there is some $p_\alpha \in G$ such that $p_\alpha \Vdash \varphi_\alpha$ or $p_\alpha \Vdash \neg\varphi_\alpha$. For every ordinal α , the formula φ_α satisfies the requirement of Lemma 22. Therefore, $p_\alpha^A \Vdash \varphi_\alpha$ if and only if $p_\alpha \Vdash \varphi_\alpha$, and hence $X \in M[G^A]$. q.e.d.

Theorem 24. In N , the cardinal κ is $\bar{\tau}$ -inaccessible.

Proof. Since our partial order has the c.c.c., it preserves cofinalities. Therefore κ is regular in $M[G]$ and therefore still regular in N . It remains to show that κ can't be embedded in any 2^λ for $\lambda < \kappa$.

Assume towards a contradiction that there is in N an injection $h : \kappa \rightarrow 2^\lambda$ for some $\lambda < \kappa$. Such an injection can be viewed as a subset of $\kappa \times \lambda$, and therefore as a set of ordinals. We apply Lemma 23 to h and get that there is some finite $A \subseteq \kappa \times \omega$ such that $h \in M[G^A]$ and thus

$$M[G^A] \models \kappa \leq 2^\lambda.$$

Keeping in mind that $\text{Card}(\mathbb{P}^A) = \aleph_0$, we can apply the proof of Lemma 15 and get that $M \models \kappa \leq 2^\lambda$ which is false. q.e.d.

Theorem 25. If $\text{Cons}(\text{ZFC} + \text{IC})$, then $\text{Cons}(\text{ZF} + \text{IC}_{\bar{\tau}/\bar{s}})$.

REFERENCES

- [Be81] Howard S. **Becker**, AD and the supercompactness of \aleph_1 , **Journal of Symbolic Logic** 46 (1981), p. 822-842
- [Bo02] Stefan **Bold**, AD und Superkompaktheit, Diplomarbeit, Rheinische Friedrich-Wilhelms-Universität Bonn, April 2002
- [Ka03] Akihiro **Kanamori**, The Higher Infinite, Springer 2003 [Springer Monographs in Mathematics]
- [Ke85] Alexander S. **Kechris**, Determinacy and the structure of $\mathbf{L}(\mathbb{R})$, in: Anil Nerode, Richard A. Shore (eds.), Recursion theory, Proceedings of the AMS-ASL summer institute held in Ithaca, N.Y., June 28–July 16, 1982, American Mathematical Society, 1985 [Proceedings of Symposia in Pure Mathematics 42], p. 271-283
- [Je03] Thomas **Jech**, Set Theory, The Third Millenium Edition, revised and expanded, Springer 2003 [Springer Monographs in Mathematics]
- [HoRu98] Paul **Howard**, Jean E. **Rubin**, Consequences of the Axiom of Choice, American Mathematical Society 1998 [Mathematical Surveys and Monographs 59]
- [FeLé63] Solomon **Feferman**, Azriel **Lévy**, Independence results in set theory by Cohen's method II (abstract), **Notices of the American Mathematical Society** 10 (1963), p. 593
- [ScZe01] Ernest **Schimmerling**, Martin **Zeman**, Square in Core Models, **Bulletin of Symbolic Logic** 7 (2001), p. 305-314

(A. Blass) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR
MI 48109-1043, U.S.A.

(I. M. Dimitriou & B. Löwe) MATHEMATISCHES INSTITUT, RHEINISCHE FRIEDRICH-WIL-
HELMS-UNIVERSITÄT BONN, BERINGSTRASSE 1, 53115 BONN, GERMANY

(B. Löwe) INSTITUTE FOR LOGIC, LANGUAGE AND COMPUTATION, UNIVERSITEIT VAN
AMSTERDAM, PLANTAGE MUIDERGRACHT 24, 1018 TV AMSTERDAM, THE NETHER-
LANDS

(B. Löwe) DEPARTMENT MATHEMATIK, UNIVERSITÄT HAMBURG, BUNDESSTRASSE 55,
20146 HAMBURG, GERMANY

E-mail address: `ablass@umich.edu`

E-mail address: `dimitri@math.uni-bonn.de`

E-mail address: `bloewe@science.uva.nl`