

New nonlinear hyperbolic groups

Richard D. Canary
University of Michigan
canary@umich.edu

Matthew Stover
Temple University
mstover@temple.edu

Konstantinos Tsouvalas
University of Michigan
tsouvkon@umich.edu

January 23, 2019

Abstract

We construct nonlinear hyperbolic groups which are large, torsion-free, one-ended, and admit a finite $K(\pi, 1)$. Our examples are built from superrigid cocompact rank one lattices via amalgamated free products and HNN extensions.

1 Introduction

In this note, we construct new examples of nonlinear hyperbolic groups. For us, a group is “nonlinear” if it does not admit a faithful representation into $GL_n(F)$ for F any field. As with previous constructions, our groups are built from superrigid cocompact lattices in rank 1 Lie groups. Previous examples were quotients of such lattices, small cancellation theory was used to show that the quotients are hyperbolic, and superrigidity results were used to see that they are nonlinear (see M. Kapovich [K05, §8]). Our construction involves simple HNN extensions and free products with amalgamation, and one can prove that the resulting groups are hyperbolic using the Bestvina–Feighn Combination Theorem [BF]. Our examples are large (i.e., have finite index subgroups that surject a free group of rank two), torsion-free, one-ended, and admit a finite $K(\pi, 1)$.

Theorem 1.1. *For any $n \geq 0$, there exist large, torsion-free, one-ended, nonlinear hyperbolic groups that admit a finite $K(\pi, 1)$, have first betti number n , and surject a free group of rank n .*

We present two related constructions, both of which begin with a cocompact torsion-free lattice Γ in $Sp(m, 1)$ (always with $m \geq 2$) or $F_4^{(-20)}$. As in M. Kapovich [K05], our proofs rely crucially on Corlette’s [Cor] and Gromov–Schoen’s [GS] generalizations of the Margulis superrigidity theorem to lattices in these groups. In what follows, let G be $Sp(m, 1)$ or $F_4^{(-20)}$ and X be the associated rank one symmetric space, i.e., quaternionic hyperbolic m -space or the Cayley hyperbolic plane.

In our first construction, we choose elements γ_1 and γ_2 of Γ associated with primitive closed geodesics of different length in the locally symmetric space X/Γ . We consider the group Λ_1 obtained by taking the HNN extension of Γ such that the stable letter conjugates γ_1 to γ_2 , i.e.,

$$\Lambda_1 = \langle \Gamma, t \mid t\gamma_1 t^{-1} = \gamma_2 \rangle.$$

We use superrigidity results to show that if Λ_1 is linear, then it admits a faithful representation ρ into $GL_n(\mathbb{R})$ and there is a totally geodesic embedding of X into the symmetric space Y_n of $GL_n(\mathbb{R})$ which

is equivariant with respect to the restriction $\rho|_\Gamma$ of ρ to Γ . Since the translation lengths of $\rho(\gamma_1)$ and $\rho(\gamma_2)$ agree in Y_n and f is totally geodesic, the translation lengths of γ_1 and γ_2 on X agree, which gives a contradiction. It follows that Λ_1 is nonlinear. The Bestvina–Feighn combination theorem [BF] implies that Λ_1 is hyperbolic, and it is clear that Λ_1 has first betti number 1, has the same cohomological dimension as Γ , admits a finite $K(\pi, 1)$, and is torsion-free. (In order to easily guarantee that Λ_1 is large, we will choose γ_1 and γ_2 to be elements of a normal, finite index subgroup of Γ of index at least 3.) We will see that it is easy to iterate this construction to produce examples with arbitrarily large first betti number.

Our second construction involves amalgamated free products and produces examples with first betti number zero. Let $\Delta = \langle \alpha, \beta \rangle$ be a malnormal, infinite index subgroup of Γ freely generated by α and β . Let $\phi : \Delta \rightarrow \Delta$ be an isomorphism such that the ratio of the translation lengths of α and β is different than the ratio of the translation lengths of $\phi(\alpha)$ and $\phi(\beta)$. We then construct

$$\Lambda_0 = \Gamma *_\phi \Gamma$$

from two copies of Γ by identifying Δ in the first copy with Δ in the second copy via the isomorphism ϕ . We argue, as before, that if Λ_0 is linear, then there is a representation ρ of Λ_0 into $\mathrm{GL}_n(\mathbb{R})$ such that the restriction of ρ to each factor determines an equivariant totally geodesic embedding of X into Y_n . It follows that the ratio of the translation lengths of α and β agrees with the ratios of the translation lengths of $\phi(\alpha)$ and $\phi(\beta)$, which we have disallowed. (In order to establish that Λ_0 is large we will also assume that Δ is contained in a normal subgroup of Γ of finite index at least 3.)

We regard the main advantage of our new constructions to be their relative simplicity and flexibility. For example, if one were given an explicit presentation of a superrigid lattice, one could easily write down an explicit presentation of a group of the form Λ_1 .

The first published examples of nonlinear hyperbolic groups are due to M. Kapovich [K05]. Gromov [Gr] used small cancellation theory to show that suitable quotients of a lattice Γ as above are infinite hyperbolic groups (see also [Ch, D, Ol]), and then Kapovich used superrigidity results to show that any linear representation of these quotients has finite image. In particular, these examples have Property (T), since they are quotients of Property (T) groups. It follows that these groups are not large and hence are not abstractly commensurable with our examples.

The paper is organized as follows. In §2, we give the details of our constructions and show that our groups have the claimed group-theoretic properties. In §3 we recall the necessary consequences of superrigidity for lattices in $\mathrm{Sp}(m, 1)$, $m \geq 2$, or $\mathrm{F}_4^{(-20)}$. The proofs of nonlinearity are given in §4.

Acknowledgments The authors are grateful to David Fisher for conversations about superrigidity, to Daniel Groves, Jason Manning, and Henry Wilton for conversations about hyperbolic groups, and to Jack Button for comments on an early draft of the manuscript. Canary and Tsouvalas were partially supported by NSF grants DMS-1306992 and DMS-1564362. Stover was partially supported by the National Science Foundation under Grant Number NSF 1361000 and Grant Number 523197 from the Simons Foundation/SFARI. The authors acknowledge support from U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 “RNMS: GEometric structures And Representation varieties” (the GEAR Network).

2 The constructions

In this section, we give the details of the constructions described in the introduction and establish the group-theoretic properties claimed there. Throughout this paper G will be either $\mathrm{Sp}(m, 1)$ for $m \geq 2$

or $F_4^{(-20)}$, so G acts by isometries on a rank one symmetric space X , which is quaternionic hyperbolic m -space or the Cayley hyperbolic plane, respectively. Then Γ will always denote a torsion-free cocompact lattice in G . In particular, Γ is hyperbolic, admits a finite $K(\pi, 1)$, $H^1(\Gamma, \mathbb{R}) = 0$, and the cohomological dimension of Γ is the dimension of X .

We first construct the examples with nontrivial first betti number. If $n \geq 2$, let $\{\gamma_1, \dots, \gamma_{2n}\}$ be primitive elements of Γ with distinct translation lengths. The associated geodesics in X/Γ are distinct, so no nontrivial power of γ_i is conjugate to a power of γ_j for $i \neq j$. We define

$$\Lambda_n = \langle \Gamma, t_1, \dots, t_n \mid t_i \gamma_i t_i^{-1} = \gamma_{i+n} \rangle.$$

to be obtained by repeated HNN extensions.

In order to construct examples which are large and have betti number zero and one, we observe that Γ contains a free, quasiconvex, malnormal subgroup Δ of rank two so that Δ is contained in a finite index, normal subgroup N of Γ of index at least three. We first note that, since Γ is residually finite, it contains a finite index, normal subgroup N of index at least three. I. Kapovich [K99, Thm 6.7] showed that every non-elementary hyperbolic group contains a malnormal quasiconvex subgroup which is free of rank two. Let F be a free malnormal quasiconvex subgroup of Γ of rank two and let D be a subgroup of $F \cap N$ which is free of rank two. Since every finitely generated subgroup of a free group is quasiconvex and F is quasiconvex in Γ , we see that D is quasiconvex in Γ . Kapovich's proof actually first constructs a free quasiconvex subgroup of rank two and then shows that this subgroup contains a free subgroup of rank two which is malnormal in the entire group. Therefore, D , and hence N , contains a subgroup Δ which is free of rank two and malnormal and quasiconvex in Γ .

Let γ_1 and γ_2 be generators of Δ with distinct translation length. Since Δ is malnormal in Γ , no nontrivial power of γ_1 is conjugate to a power of γ_2 . Let Λ_1 be the HNN extension of Γ given by

$$\Lambda_1 = \langle \Gamma, t \mid t \gamma_1 t^{-1} = \gamma_2 \rangle.$$

(If we do not require Λ_1 to be large, it would suffice to choose γ_1 and γ_2 to be primitive elements with distinct translation length as in the construction of Λ_n when $n \geq 2$.)

We now construct the examples with trivial first betti number. Let α and β generate Δ and let $\phi : \Delta \rightarrow \Delta$ be an isomorphism such that the ratio of the translation lengths of α and β is different than the ratio of the translation lengths of $\phi(\alpha)$ and $\phi(\beta)$. We define

$$\Lambda_0 = \Gamma *_\phi \Gamma.$$

to be obtained from two copies of Γ by identifying Δ in the first copy with Δ in the second copy via the isomorphism ϕ . (If we do not require that Λ_0 is large, it would suffice to choose Δ to be the malnormal, quasiconvex subgroup of Γ guaranteed by Kapovich [K99].)

Proposition 2.1. *For all n , a group Λ_n constructed as above is hyperbolic, torsion-free, large, one-ended, has a finite $K(\pi, 1)$, has first betti number n , and its cohomological dimension is the dimension of X . Moreover, if $n \geq 1$, Λ_n admits a surjective homomorphism to the free group F_n of rank n .*

Proof. That Λ_n is torsion-free, one-ended, has a finite $K(\pi, 1)$, has first betti number n , and has cohomological dimension equal to the dimension of X follows from standard facts about graphs of groups (see, for example, Serre [Ser, Chap. 1] or Scott–Wall [SW]). If $n \geq 1$, then Λ_n clearly surjects onto the group freely generated by $\{t_1, \dots, t_n\}$. The fact that each Λ_n is hyperbolic is a special case of the Bestvina–Feighn combination theorem [BF], which is explicitly stated in I. Kapovich [K01, Ex. 1.3] as follows:

Theorem 2.2.

1. If A and B are hyperbolic groups and C is a quasiconvex subgroup of both A and B that is malnormal in either A or B , then $A *_C B$ is hyperbolic.
2. If A is a hyperbolic group and a_1 and a_2 are elements of A so that no nontrivial power of a_1 is conjugate to a power of a_2 , the HNN extension

$$\langle A, t \mid ta_1t^{-1} = a_2 \rangle$$

is hyperbolic.

Part (1) immediately implies that Λ_0 is hyperbolic, while part (2) gives that Λ_n is hyperbolic if $n \geq 1$. Also, notice that normal form for words in the HNN extension Λ_{n-1} (see [Ser, §I.5]) implies we still have that no power of γ_n is conjugate to a power of γ_{2n} in Λ_{n-1} .

We remarked above that Λ_n is large for $n \geq 2$, so it remains to prove that Λ_1 and Λ_0 are also large. Suppose that $n = 1$. There is a surjective homomorphism

$$p_1 : \Lambda_1 \rightarrow H_1 = \Gamma/N * \mathbb{Z}$$

given by projecting Γ onto Γ/N and taking t to the generator of \mathbb{Z} . Let J be a finite index subgroup of H_1 which is isomorphic to a free group of rank at least two, which exists, since Γ/N has order at least three. Then $p_1^{-1}(J)$ is a finite index subgroup of Λ_1 and p_1 restricts to a surjection of $p_1^{-1}(J)$ onto J , so Λ_1 is large.

We now consider Λ_0 . There exists a surjective homomorphism

$$p_0 : \Lambda_0 \rightarrow H_0 = \Gamma/N * \Gamma/N$$

given by projecting the first factor of Λ_0 to the first factor of H_0 and the second factor of Λ_0 to the second factor of H_0 . Notice that this is well-defined since Δ has trivial image in both factors. As above, H_0 contains a finite index subgroup which is isomorphic to a free group of rank at least two, so Λ_0 is large. \square

Remarks: 1) Kapovich [K99] further uses a malnormal quasiconvex free subgroup of a word hyperbolic group G to construct a hyperbolic group G^* which contains G as a non-quasiconvex subgroup. We note that G^* is a quotient of a group of the form Λ_2 , obtained by identifying the two stable letters, so if G is a superrigid rank one lattice then G^* can be chosen to be nonlinear.

2) We expect that the techniques of Belegradek–Osin [BO], which also begin with quotients of superrigid lattices and employ more powerful small cancellation theoretic results, also produce large, one-ended, nonlinear hyperbolic groups (in particular, see [BO, Thm. 3.1]).

3) It is clear that one can construct infinitely many isomorphism classes of groups of the form Λ_n , for each n , even if one begins with a fixed superrigid lattice Γ . For example, if $n \geq 1$, it follows readily from the JSJ theory for hyperbolic groups, see Sela [Sel], that the isomorphism type of a group of the form Λ_1 is determined, up to finite ambiguity, by the conjugacy class of the pair $\{\gamma_1, \gamma_2\}$ in Γ .

3 Superrigidity

In this section, we record a version of the superrigidity theorem of Corlette [Cor] and Gromov–Schoen [GS] that is crafted for our purposes. In our statement Y_n will denote the symmetric space

$$Y_n = ZO(n) \backslash GL_n(\mathbb{R}) = PO(n) \backslash PGL_n(\mathbb{R})$$

associated with $GL_n(\mathbb{R})$, where Z denotes the center of $GL_n(\mathbb{R})$.

Theorem 3.1. *Suppose that Γ is a lattice in G , where G is either $Sp(m, 1)$ or $F_4^{(-20)}$, F is a field of characteristic zero, and $\rho : \Gamma \rightarrow GL_d(F)$ is a representation with infinite image.*

1. *There exists a faithful representation $\tau : GL_d(F) \rightarrow GL_n(\mathbb{R})$ for some n such that $\tau \circ \rho(\Gamma)$ has noncompact Zariski closure.*
2. *If $F = \mathbb{R}$ and $\rho(\Gamma)$ has noncompact Zariski closure in $GL_d(\mathbb{R})$, then there exists a ρ -equivariant totally geodesic map*

$$f_\rho : X \rightarrow Y_d,$$

where $X = K \backslash G$ is the symmetric space associated with G .

Proof. Since Γ is finitely generated we may assume that F is isomorphic to a subfield of \mathbb{C} . Moreover, $GL_d(\mathbb{C})$ is a subgroup of $GL_{2d}(\mathbb{R})$. It follows that there exists an injective representation $\eta : GL_d(F) \rightarrow GL_n(\mathbb{R})$ for some n , so we may assume that the original representation maps into $GL_n(\mathbb{R})$.

Fisher and Hitchman [FH, Thm. 3.7] then observe that the existing results on superrigidity imply that one can factor ρ as two representations

$$\rho_i : \Gamma \rightarrow GL_{n_i}(\mathbb{R}) \subseteq GL_n(\mathbb{R})$$

such that:

1. When ρ_1 is nontrivial, there is a group G' locally isomorphic to G , a continuous representation $\hat{\rho}_1 : G' \rightarrow GL_{n_1}(\mathbb{R})$, and an embedding $\iota : \Gamma \hookrightarrow G'$ of Γ as a lattice in G' such that $\rho_1 = \hat{\rho}_1 \circ \iota$.
2. The image of ρ_2 is bounded, i.e., has compact Zariski closure.
3. The groups $\rho_1(\Gamma)$ and $\rho_2(\Gamma)$ commute and $\rho(\gamma) = \rho_1(\gamma)\rho_2(\gamma)$ for all $\gamma \in \Gamma$.

If ρ_1 is nontrivial, the continuous embedding $\hat{\rho}_1 : G' \rightarrow GL_{n_1}(\mathbb{R})$ determines a totally geodesic embedding of X into Y_{n_1} , hence into Y_n . Since ρ_1 and ρ_2 commute, this is a ρ -equivariant map.

When ρ_1 is trivial, we follow arguments in the proof of [K05, Thm. 8.1]. Note that our use of [FH, Thm. 3.7] allows us to know beforehand that the solvable radical considered in [K05] is trivial. As in [K05], the fact that Γ has Property (T) allows us to conclude that we may conjugate ρ so that $\rho(\Gamma) \subseteq GL_n(k)$ for some number field k . Given an element $\sigma \in \text{Aut}(k/\mathbb{Q})$, we can choose an extension of σ to an element of $\text{Aut}(\mathbb{C}/\mathbb{Q})$, which we continue to denote by σ . Applying σ to matrix entries induces an embedding $\tau_\sigma : GL_n(F) \rightarrow GL_n(\mathbb{C})$.

Following the adelic argument in [K05], if $\rho(\Gamma)$ were bounded for every valuation of k then $\rho(\Gamma)$ would be finite, which is a contradiction. Moreover, $\rho(\Gamma)$ must be bounded for every nonarchimedean valuation by nonarchimedean superrigidity [GS]. Consequently, there exists $\sigma \in \text{Aut}(k/\mathbb{Q})$ such that $\tau_\sigma(\rho(\Gamma))$ has noncompact Zariski closure in $GL_n(\mathbb{R})$ or $GL_{2n}(\mathbb{R})$, according to whether $\sigma(k) \otimes_{\sigma} \mathbb{R}$ is \mathbb{R} or \mathbb{C} . Applying the previous argument to $\tau_\sigma \circ \rho$, there is a $(\tau_\sigma \circ \rho)$ -equivariant totally geodesic embedding of X into Y_n or Y_{2n} , accordingly. This completes the sketch of the proof. \square

M. Kapovich [K05] also points out that superrigidity rules out faithful representations of Γ into linear groups of fields of positive characteristic. Briefly, one shows that the image of ρ lies in $\mathrm{GL}_n(k)$ where k is a finite extension of $\mathbb{F}_p(x_1, \dots, x_n)$. Then, applying Gromov–Schoen superrigidity [GS] to each valuation of k associated with some $x_i^{\pm 1}$, one sees that $\rho(\Gamma)$ is bounded in each field associated with such a valuation on k , as all valuations on k are nonarchimedean. It follows that $\rho(\Gamma)$ is bounded and hence finite. Thus we have:

Proposition 3.2. *If Γ is a lattice in either $Sp(m, 1)$ or $F_4^{(-20)}$ and F is a field of characteristic $p > 0$, then there does not exist a faithful representation of Γ into $\mathrm{GL}_n(F)$ for any n .*

4 Proofs of nonlinearity

To complete the proof of Theorem 1.1 it remains to prove:

Theorem 4.1. *Groups of the form Λ_n constructed in Section 2 are nonlinear.*

Proof. We begin with a group of the form

$$\Lambda_1 = \langle \Gamma, t \mid t\gamma_1 t^{-1} = \gamma_2 \rangle$$

constructed in Section 2, where Γ is a cocompact lattice in G and G is either $Sp(m, 1)$ or $F_4^{(-20)}$. Recall that X is the symmetric space associated with G and that γ_1 and γ_2 are assumed to have different translation lengths on X .

Suppose that F is a field and $\eta : \Lambda_1 \rightarrow \mathrm{GL}_d(F)$ is a faithful representation. Applying Proposition 3.2 to the restriction $\rho = \eta|_{\Gamma}$ of η to Γ , we conclude that F has characteristic zero. Theorem 3.1 implies that there exists a faithful representation $\tau_{\sigma} : \mathrm{GL}_d(F) \rightarrow \mathrm{GL}_n(\mathbb{R})$, for some n and a $(\tau_{\sigma} \circ \rho)$ -equivariant totally geodesic embedding f of X into Y_n , where Y_n is the symmetric space associated with $\mathrm{GL}_n(\mathbb{R})$.

Since $\tau_{\sigma}(\rho(\gamma_1))$ is conjugate to $\tau_{\sigma}(\rho(\gamma_2))$ in $\tau_{\sigma}(\eta(\Lambda_1))$, and hence in $\mathrm{GL}_n(\mathbb{R})$, they have the same translation length on Y_n . However, since f is a $(\tau_{\sigma} \circ \rho)$ -equivariant totally geodesic embedding, this implies that γ_1 and γ_2 have the same translation length in X , which is a contradiction, hence Λ_1 is nonlinear. Notice that if $n \geq 2$, then any group of the form Λ_n constructed in Section 2 contains a subgroup of the form Λ_1 , so Λ_n is also nonlinear.

Now suppose we have a group of the form

$$\Lambda_0 = \langle \Gamma_1, \Gamma_2 \mid \alpha_1 = \phi(\alpha)_2, \beta_1 = \phi(\beta)_2 \rangle$$

where each Γ_i is a copy of Γ , $\Delta = \langle \alpha, \beta \rangle$ is a subgroup of Γ freely generated by α and β , Δ_i is the copy of Δ in Γ_i and if $\delta \in \Delta$, then δ_i is the copy of δ in Δ_i . Moreover, ϕ is an automorphism of Δ so that the ratio of the translation lengths of α and β on X differs from the ratio of translation lengths of $\phi(\alpha)$ and $\phi(\beta)$ on X .

Suppose that F is a field and $\eta : \Lambda_0 \rightarrow \mathrm{GL}_d(F)$ is a faithful representation. Let $\rho_1 = \eta|_{\Gamma_1}$ and $\rho_2 = \eta|_{\Gamma_2}$. We again apply Proposition 3.2 to conclude that F has characteristic zero, Theorem 3.1 implies that there exists a faithful representation $\tau_{\sigma} : \mathrm{GL}_d(F) \rightarrow \mathrm{GL}_n(\mathbb{R})$, for some n and a $(\tau_{\sigma} \circ \rho_1)$ -equivariant embedding f of X into Y_n , where Y_n is the symmetric space associated with $\mathrm{GL}_n(\mathbb{R})$. Since $\tau_{\sigma}(\rho_1(\Delta_1)) = \tau_{\sigma}(\rho_2(\Delta_2))$ has noncompact Zariski closure, Theorem 3.1 implies that there exists a $(\tau_{\sigma} \circ \rho_2)$ -equivariant embedding g of X into Y_n . Notice that $\tau_{\sigma}(\rho_1(\alpha_1)) = \tau_{\sigma}(\rho_2(\phi(\alpha)_2))$ and that $\tau_{\sigma}(\rho_1(\beta_1)) = \tau_{\sigma}(\rho_2(\phi(\beta)_2))$.

Since f and g are equivariant totally geodesic embeddings, there exist positive constants c_1 and c_2 so that if $\gamma \in \Gamma$, then the ratio of the translation length of $\tau_\sigma(\rho_i(\gamma_i))$ on Y_n and the translation length of γ on X is c_i . Indeed, the metrics on $f(X)$ and $g(X)$ differ by a scalar multiple. It follows that the ratio of the translation lengths of α and β on X agrees with the ratio of the translation lengths of $\phi(\alpha)$ and $\phi(\beta)$ on X . However, this contradicts our assumptions, so Λ_0 is nonlinear. \square

References

- [BO] M. Belegradek and D. Osin, Rips construction and Kazhdan property (T), *Groups, Geom., Dyn.* **2** (2008) 1–12.
- [BF] M. Bestvina and M. Feighn, A combination theorem for negatively curved groups. *J. Diff. Geom.* **35** (1992), 85–101.
- [Ch] C. Champetier, Petite simplification dans les groupes hyperboliques. *Ann. Fac. Sci. Toulouse Math.* **3**(1994), 161–221.
- [Cor] K. Corlette, Archimedean superrigidity and hyperbolic geometry. *Ann. of Math. (2)* **135** (1992), 165–182.
- [D] T. Delzant, Sous-groupes distingués et quotients des groupes hyperboliques. *Duke Math. J.* **83** (1996), 661–682.
- [FH] D. Fisher and T. Hitchman, Strengthening Kazhdan’s Property (T) by Bochner methods. *Geom. Dedicata* **160** (2012), 333–364.
- [Gr] M. Gromov, Hyperbolic groups. *Essays in group theory*, 75–263, Math. Sci. Res. Inst. Publ., **8**, Springer 1987.
- [GS] M. Gromov and R. Schoen, Harmonic maps into singular spaces and p-adic superrigidity for lattices in groups of rank one. *Inst. Hautes Études Sci. Publ. Math.* **76** (1992), 165–246.
- [K99] I. Kapovich, A non-quasiconvexity embedding theorem for hyperbolic groups, *Math. Proc. Camb. Phil. Soc.* **127**(1999), 461–486.
- [K01] I. Kapovich, The combination theorem and quasiconvexity. *Int. J. Alg. Comp.* **11** (2001), 185–216.
- [K05] M. Kapovich, Representations of polygons of finite groups. *Geom. Top.* **9** (2005), 1915–1951.
- [Ol] A.Y. Ol’shanskii, On residualing homomorphisms and G -subgroups of hyperbolic groups. *Int. J. Alg. Comp.* **3**(1993), 365–409.
- [SW] P. Scott and T. Wall, Topological methods in group theory. *Homological Group Theory*, L.M.S. Lecture Note Series vol. **36**, Camb, Univ. Press, 1979, 137–203.
- [Sel] Z. Sela, Structure and rigidity in (Gromov) hyperbolic groups and discrete groups in rank 1 Lie groups. II. *G.A.F.A.* **7**(1997), 561–593.
- [Ser] J.-P. Serre, *Trees*. Springer Monographs in Mathematics. Springer-Verlag, 1980.