

Equivariant and Non-equivariant homotopy Theory IV

Note Title

1/19/2012

Adams S.S for spheres

$$E_2 = E_{x+}^{s,t}(\mathbb{Z}/2, \mathbb{Z}/2) \Rightarrow \pi_* S^0$$

Kervaire Inv. 1 manifold

\leftrightarrow elts. represented in E_2

by $h_i \in E_{x+}^{2^i, 2^{i+1}}$

Hill-Hopkins-Ravenel (2009):

h_i^2 do not survive to E_∞
for $i \geq 7$

i.e. no Kervaire Inv. 1 manifold
of dim $2^{n+1}-2$ for $n \geq 7$

General plan:

construct a spectrum Ω

from $\mathbb{Z}/8$ -equivariant

1. take Real cobordism $M\mathbb{R}$,
4 copies

$$M\mathbb{R} \wedge M\mathbb{R} \wedge M\mathbb{R} \wedge M\mathbb{R}$$

$M\mathbb{R}^{(4)}$, $\mathbb{Z}/8$ -equiv.

2. invert a homotopy class D
of $M\mathbb{R}^{(4)} \Rightarrow D^{-1} M\mathbb{R}^{(4)}$

3. take fixed points: Ω

3 facts about Ω :

1. detection theorem: if $\theta_i \in \pi_{2^{i+1}-2} S^0$ is represented by h_i^2 , then they map to non-0 elts. in $\pi_{2^{i+1}-2} \Omega$

2. gap theorem: $\pi_{-2} \Omega = 0$

3. periodicity theorem.

$$\pi_* \Omega \cong \pi_{*+256} \Omega$$


these three give Kervaire Inv. 1
Th.

Main Tool: slice Spectral
sequence, from slice tower

analogue of Postnikov tower:

given X , "kill" all homotopy
classes $S^k \rightarrow X$ for $k > n$

at n -th stage



attach cones
 on all $S^k \rightarrow X$,
 $k > n$

slice tower: equivariant analogue

replace S^k by "slice cells"

$$G \ltimes_{\rho_K} S^m$$

$K \subseteq G$ subgroup, $m \in \mathbb{Z}$,

$\rho_K =$ regular rep. of K

also allow Σ^{-1} of these

for X equivariant:

$$\dots \longrightarrow P^n X \longrightarrow P^{n-1} X \longrightarrow \dots$$

$$\begin{array}{c} \uparrow \\ P_n^n X = \text{homotopy fiber} \\ = n\text{-th slice of } X \end{array}$$

long exact sequences in
homotopy groups

\Rightarrow slice S.S., $RO(G)$ -graded

easier to write each "twist"
separately; given virtual rep. V

$$\mathbb{E}_2^{s,t} = \pi_{V+t-s} P_{|V|+t} X \Rightarrow \pi_{V+t-s} X$$

H-H-R: for MTR , $MTR^{(2)}$, $MTR^{(4)}$

computed what the slices are,
homotopy gps.

for $X = M\mathbb{R}^{(2^{n-1})}$, $\mathbb{Z}/2^n$ -equivariant

$$n = \text{odd}: P_n^n X = *$$

$$n = \text{even}: P_n^n X = \left(\bigvee_{\substack{\text{slice cells} \\ \text{of dim } n}} \right) \wedge H\underline{\mathbb{Z}} \\ K \neq \{e\}$$

Gap th: in this case,

$$\pi_*^G (\text{slice cell} \wedge H\underline{\mathbb{Z}})_i = 0 \quad \text{in} \\ \text{dim } -4 < i < 0$$

Periodicity Th: look at differentials

(we did for MTR)

Igor's talk: $d_{2^{k+1}-1}(\sigma^{-2^k}) = v_k a^{2^{k+1}-1}$

$\alpha = \text{sign rep.}$

$$|a| = -\alpha, \quad |\sigma| = \alpha - 1, \quad |v_k| = (2^k - 1)(1 + \alpha)$$

similar elts. in slice S.S.

E_2 -term for $MTR^{(2^{n-1})}$

1. $a \in E_2^{1, 1-\alpha}$, $\alpha = \text{sign rep. of } \mathbb{Z}/(2^n)$

2. $\sigma^{-2} \in E_2^{0, 2-2\alpha}$

(any oriented rep. V , get

$$u_V \in E_2^{0, |V|-\dot{V}}$$


3. $f_i \in E_2^{(g-1)i, gi}$

$f_{2^k-1} = V_k a^{2^k-1}$

$V = -2^k \alpha$

$g = |G|$

slope g^{-1}



$$d_{1+(2^k-1)g}(\sigma^{-2^k}) = a^{2^k} f_{2^k-1}$$

$$= a^{2^{k+1}-1} v_k$$

Idea: invert certain multiples
of v_k , will make powers of
 σ^{-2^k} into permanent cycles
once we can divide by these
 v_k 's, target of differentials
from σ^{-2^k} will be killed by

by earlier differentials

need a product of several factors, from MTR , $MTR^{(2)}$, $MTR^{(4)}$

$$D \in \pi_{19\rho_8} MTR^{(4)}$$

invert $D \Rightarrow u_{32\rho_8}$ is perm. cycle

multiply by a factor of D

to put it into integral dim

becomes a unit in

$$\pi_{256}^G D^{-1} MTR^{(4)}$$

Detection Th. use Adams-Novikov

S.S., another tool for $\pi_* S^0$:
use MU instead of $H\mathbb{Z}/2$

$$\text{Ext}_{\text{MU}_* \text{MU}}^{\text{MU}_*}(\text{MU}_*, \text{MU}_*) \Rightarrow \pi_* S^0$$

$\text{ANSS for } S^0 \rightarrow \text{ANSS for } \Omega \rightarrow \text{homotopy fixed pt. SS. for } D^{-1} \text{MTR}^{(4)}$
 \downarrow
 $\text{ASS for } S^0$
 h_i^2

point: if x in ANSS in $S^0 \oplus$
 goes to h_i^2 , it maps to
 non-0 in homotopy fixed
 pt. S.S. for $D^{-1} \text{MTR}^{(4)}$

define

$$R_* = \mathbb{Z}_2[\xi_8][w^{\pm 1}]$$

← 8-th root of 1

$$H^*(\mathbb{Z}/8, D^{-1}MR_*^{(4)}) \rightarrow H^*(\mathbb{Z}/8, R_*)$$

⊕
computable