## $\rm hw10$ , due: Wednesday, November 28

1. The age of organic remains is determined by <u>radiocarbon dating</u> as follows. When cosmic rays enter the atmosphere, they convert nitrogen to a radioactive isotope of carbon, <sup>14</sup>C, with a half-life of 5730 years. Living animals absorb <sup>14</sup>C in the atmosphere and food chain, and when they die, the <sup>14</sup>C present in the remains decreases by radioactive decay. Consider the case of parchment, a thin sheet made from animal skins used for writing in ancient times. A parchment fragment was discovered having 74% as much <sup>14</sup>C as in current living animal skins. Find the age of the fragment.

2. Find the sum of the series  $\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \frac{1}{2} + \frac{1}{6} + \frac{1}{12} + \cdots$  two ways.

method a) Express  $\frac{1}{n(n+1)}$  by partial fractions; examine the partial sums  $s_n$  of the series. method b) Sketch the curves  $y = x^n$  for  $0 \le x \le 1$  and n = 0, 1, 2, 3, 4 on a common plot; let  $A_n$  be the area between  $y = x^n$  and  $y = x^{n+1}$  for any  $n \ge 0$ ; compute  $A_n$  in terms of n; express the area of the unit square in terms of the  $A_n$ ; deduce the sum of the series.

3. Find the interval of convergence and sum of the power series.

a) 
$$\sum_{n=0}^{\infty} (x-4)^n$$
 b)  $\sum_{n=0}^{\infty} \frac{(-1)^n x^n}{n+1}$  c)  $\sum_{n=1}^{\infty} \frac{nx^n}{3^n}$ 

4. Three infinite sequences of tangent circles approach the vertices of an equilateral triangle. The figure shows the first few circles. Assume the triangle has sides of length 1. Express the area covered by the circles as a series. What fraction of the triangle area is covered by the circles?



5. This problem continues the discussion of Taylor approximation. Given a function f(x) and a point x = a, define  $T_3(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3$ ; it is a cubic function of x called the <u>Taylor polynomial of degree 3</u>.

a) Show that  $T_3^{(n)}(a) = f^{(n)}(a)$  for n = 0, 1, 2, 3, where the superscript denotes the *n*th derivative; thus  $T_3(x)$  and f(x) agree to 3rd order at x = a, and  $T_3(x)$  is a <u>cubic approximation</u> to f(x). b) Recall from hw9:  $f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2}(x-a)^2 + \int_a^x \frac{(x-t)^2}{2}f'''(t) dt$ . Now show that  $f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \int_a^x \frac{(x-t)^3}{3!}f^{(4)}(t) dt$ . (hint: in the integral from the hw9 result, set u = f'''(t),  $dv = \frac{(x-t)^2}{2}dt$ , and integrate by parts) c) Using part (b), derive the error bound  $|f(x) - T_3(x)| \le \frac{1}{4!}M_4|x-a|^4$ , where  $M_4 = \max |f^{(4)}(t)|$ . (hint: part (b)  $\Rightarrow f(x) = T_3(x) + \int_a^x \frac{(x-t)^3}{3!}f^{(4)}(t) dt$ ; then follow the steps from hw7, problem 10) d) Let  $f(x) = e^x$ , a = 0. Find  $T_3(x)$ . Sketch  $f(x), T_3(x)$  on the same graph around x = a. e) Make a table with the following format. column 1: |x - a|, column 2:  $|f(x) - T_3(x)|$ . Take  $f(x) = e^x, a = 0$  and fill in the entries for x = 1, 1/2, 1/4, 1/8 using a calculator. When |x - a|is reduced by a factor of  $\frac{1}{2}$ , by what factor is the error  $|f(x) - T_3(x)|$  reduced?