

Scott complexity and adjoining roots to finitely generated groups

Larsen Louder

August 16, 2011

Abstract

We give several generalizations of the well known fact that a commutator in a free group is not a proper power.

1 Introduction

We consider the problem of adjoining roots to elements of finitely generated groups. An element of a free group is primitive if it is part of a basis. The letter \mathbb{F} will be used to represent a nonabelian free group. Let G be a group. An element $g \in G$ is *indivisible* if isn't a proper power. The group $\langle G, \gamma'_i | (\gamma'_i)^{k_i} \gamma_i^{-1}, i \dots n \rangle$ is denoted $G[\sqrt[k_i]{\gamma_i}]$.

By [She55], if an amalgam $A_1 *_{\langle t \rangle} A_2$ is finitely generated and free, then t is primitive in one of A_1 or A_2 . If $G[\sqrt[k]{\gamma}]$ is free, it corresponds to the case $A_2 \cong \mathbb{Z}$, and since γ is not primitive in $\langle \sqrt[k]{\gamma} \rangle$, γ is primitive in G . The identity map $A_1 *_{\langle t \rangle} A_2 \rightarrow A_1 *_{\langle t \rangle} A_2$ is a morphism of free groups, and the purpose of this paper is to examine what happens when we are simply given a map $A_1 *_{\langle t \rangle} A_2 \rightarrow \mathbb{F}$ embedding A_i .

Consider Vaught's equation $a^2 b^2 c^2 = 1$. Lyndon showed that if a , b , and c satisfy Vaught's equation in a free group then they commute. Let $V = \langle a, b, c \mid a^2 b^2 c^2 \rangle$. Solutions to $a^2 b^2 c^2 = 1$ in a group G correspond to elements of the set $\text{Hom}(V, G)$. The group V is the fundamental group of the connected sum of three projective planes, or equivalently, the connected sum of a torus T and a projective plane P . Lyndon's theorem simply says that all elements of $\text{Hom}(V, \mathbb{F})$ factor through the map killing P . Equivalently, a commutator in a free group isn't a square. Lyndon and Schützenberger [LS62] (See also [Sch59].) extended this fact to equations $a^p b^q c^r = 1$, $p, q, r \geq 2$, of the same type.

More generally, if (y_1, \dots, y_{n+1}) is a solution to $\delta(x_1, \dots, x_n) = x_{n+1}^k$, $k > 1$ in \mathbb{F} , and δ is neither a proper power nor primitive in $\langle x_1, \dots, x_n \rangle$, then the subgroup of \mathbb{F} generated by $\{y_1, \dots, y_{n+1}\}$ has rank less than n [Bau65]. The theorems of Lyndon, Lyndon-Schützenberger, and Baumslag may all be reinterpreted as statements about maps of graphs of free groups to free groups embedding vertex groups.

The purpose of this paper is to generalize this fact to (most) arbitrary finitely generated groups. We define a complexity which reduces to the rank in the case of a free group, behaves well with respect to adjunctions of roots to arbitrarily many elements, and strictly decreases if roots are added to (mutually) nonprimitive (collections of) elements. The following theorem is an important step in our proof that limit groups have finite Krull dimension [?].

Definition 1.1 ([Sco73, Swa04]). Let G be a finitely generated group with Grushko decomposition $G_1 * \dots * G_p * \mathbb{F}_q$. The *Scott complexity* of G is the ordered pair $\chi_{\text{fd}}(G) = (q - 1, p)$.

Theorem 1.2. *Let G be a finitely generated group, and let $\gamma_1, \dots, \gamma_n$ and $\delta_1, \dots, \delta_m$ be nontrivial collections of indivisible elements of G , such that all γ_i are contained in freely indecomposable free factors of G , no δ_j is contained in any freely indecomposable free factor of G , and all δ_j have nonconjugate centralizers. Suppose $G \hookrightarrow H$ and that the inclusion map extends to a surjection*

$$G' = G \left[\sqrt[k]{\gamma_i}, \sqrt[l]{\delta_j} \right] \twoheadrightarrow H$$

Then $\chi_{\text{fd}}(G) \geq \chi_{\text{fd}}(H)$. If equality holds and H has no \mathbb{Z}_2 free factors then there are elements d_j of G such that

$$G \cong G_1 * \dots * G_p * F * \langle \delta_1^{d_1} \rangle * \dots * \langle \delta_m^{d_m} \rangle$$

with G_i freely indecomposable, and F free.

Baumslag's theorem clearly corresponds to the case $\chi_{\text{fd}}(G) = \chi_{\text{fd}}(H) = (q - 1, 0)$, $n = 0$, $m = 1$.

Example (No \mathbb{Z}_2 free factors of H is necessary.). *Let $G = G_1 * G_2$, $H = H_1 * \mathbb{Z}_2$, where $H_1 = G_1 * G_2 / \langle\langle \alpha_1 = \alpha_2 \rangle\rangle$, $\alpha_i \neq 1 \in G_i$, and x generates \mathbb{Z}_2 . Then H is a quotient of $G \sqrt[\alpha_1 \alpha_2]$. The inclusion $G \hookrightarrow H$ maps G_1 to $G_1 < H_1$ and G_2 to $xG_2x < xH_1x$. Then $\alpha_1 \rightarrow \alpha_1$, $\alpha_2 \rightarrow x\alpha_1x$, and $\alpha_1\alpha_2 \rightarrow (\alpha_1x)^2$.*

Our analysis also gives the following.

Theorem 1.3 (Improved Shenitzer's Theorem). *Let $G = A_1 *_t A_2$, A_1 and A_2 free of rank n and m , respectively. If t is indivisible in at least one vertex group and G maps onto a free group of rank $n + m - 1$, then G is free and t is primitive in at least one A_i . If t is not indivisible in either factor then G is a free product of a free group and an amalgamated product of two infinite cyclic groups.*

Baumslag suggests a conjugacy separability problem for elements of the free group. If α and β are nonconjugate elements of \mathbb{F}_n , then either (without loss) there is a free factorization $\mathbb{F}_n \cong F * \langle \alpha' \rangle$ such that α is conjugate of a power of α' and β is conjugate into F , or $\langle \mathbb{F}_n, t | t\alpha t^{-1}\beta^{-1} \rangle$ can map onto a free group of rank at most $n - 1$.

Theorem 1.4. *Let F be a free group of rank $n > 1$, and let $\{t_i\}$ be stable letters, and let $\{\delta_j\}$ be nontrivial, nonconjugate elements of F . Let G be the group $F * \langle t_i \rangle / \langle\langle t_i \delta_{j_i} t_i^{-1} = \delta_{j'_i} \rangle\rangle$, and assume that each δ_j participates. If G has a map onto a free group of rank n embedding F , then $F \cong F' * \langle \delta_j \rangle$ for some j , with each element $\gamma_{j'}$, $j' \neq j$, conjugate (in F) into F' .*

Acknowledgments

It's hard to overstate my gratitude to Mark Feighn for his careful reading of more versions of this paper than I can count. I would also like to thank Mladen Bestvina for pointing out the proof of Shenitzer's theorem and teaching me folding.

2 Free Groups

2.1 Immersions I

A graph is a one dimensional CW complex. A graph with boundary Γ is a graph with some valence one vertices marked as elements of the boundary $\partial\Gamma$. A graph is a *core graph* if it has no valence one vertices outside its boundary. A morphism of graphs is a map that sends vertices to vertices and maps interiors of edges homeomorphically to interiors of edges. A morphism of graphs is an *immersion* if it is locally injective. Equivalently, a morphism of graphs is an immersion if it is injective on links of vertices. Stallings observed that if $f: A \hookrightarrow B$ is a morphism of finitely generated free groups then there is an immersion of finite graphs $\varphi: \Gamma_A \rightarrow \Gamma_B$ such that after identifying A and B with the fundamental groups of Γ_A and Γ_B appropriately, φ represents f .




Figure 1: A subdivided mapping cylinder $S^1 \xrightarrow{\times 3} S^1$.

If Γ is a graph and p is a closed path crossing some edge of Γ exactly once then p represents a primitive element in $\pi_1(\Gamma)$.

Let b and r be circles, and let $l: b \rightarrow r$ be the l -fold cover. The mapping cylinder $M(l)$ of l is the space $b \times [0, 1] \sqcup r/x \sim l(x)$. We call $r \subset M(l)$ the *core*, and we call $b \times \{0\} \subset M(l)$ the *boundary*. Let Γ be a graph, and let $\gamma: S^1 \rightarrow \Gamma$ be a cyclically reduced path, and suppose that γ has an l -th root, that is, a cyclically reduced path $\sqrt[l]{\gamma}: S^1 \rightarrow \Gamma$ such that $(\sqrt[l]{\gamma})^l$ is homotopic to γ through reduced edge paths. By the mapping cylinder $M(\gamma, l)$ we mean the mapping cylinder of the l -fold cover of S^1 , subdivided into rectangles, such that the edges of squares not contained in the boundary or core are sent to vertices, and the restrictions to $b \times \{x\}$ are reduced edge paths, and with the natural map $M(\gamma, l) \rightarrow \Gamma$ induced by γ , $\sqrt[l]{\gamma}$, and the homotopy. See Figure 1.

If $G = A_1 *_{\langle t \rangle} A_2$ is free, then each map $A_i \hookrightarrow G$ may be represented by an immersion $\iota_i: \Gamma_{A_i} \hookrightarrow R$, where R is a graph with fundamental group identified with G . Represent $\langle t \rangle$ by a pair of immersions $S^1 \hookrightarrow \Gamma_{A_i}$, and glue the boundary components of an (subdivided) annulus to $\Gamma_{A_1} \sqcup \Gamma_{A_2}$ along the immersions of S^1 to obtain a graph of spaces X . Extend ι_i to a map $\varphi: X \rightarrow R$. Pull back midpoints of edges of R to produce embedded graphs in X “transverse” to both Γ_1 and Γ_2 . The hypothesis that ι_i and the maps of S^1 are immersions implies that the preimage graphs must be finite forests, otherwise the identity map $G \rightarrow G$ has nontrivial kernel. This implies that the representative of t as an immersion, in at least one Γ_i , must cover some edge exactly once, and t is therefore a basis element in some A_i .

Consider Baumslag’s theorem: A solution to $\delta(x_1, \dots, x_n) = x_{n+1}^k$, δ a word in x_i and $k > 1$, in a free group \mathbb{F} corresponds to a map $\langle x_1, \dots, x_n \rangle \left[\sqrt[k]{\delta} \right] \rightarrow$

\mathbb{F} . If the image of $\langle x_1, \dots, x_n \rangle$ in \mathbb{F} has rank less than n then the conclusion clearly holds, so we may assume that the map is an embedding on $\langle x_i \rangle$. The group $\langle x_1, \dots, x_n \rangle \left[\sqrt[k]{\delta} \right]$ is an amalgam of a free group and a cyclic group. We construct a space X and a map $X \rightarrow R$ representing the map $\langle x_1, \dots, x_n \rangle \left[\sqrt[k]{\delta} \right] \rightarrow \mathbb{F}$ exactly as in the proof of Shenitzer's theorem. The main observation of this paper is that if the image of $\langle x_1, \dots, x_n \rangle \left[\sqrt[k]{\delta} \right]$ has rank n then the preimage graphs are forestlike, and if $k > 0$ and δ is indivisible, then they are forests. Using this and the fact that a closed path crossing some edge only once represents a primitive element we conclude that δ must be primitive. In the following subsection we make this argument precise.

2.2 Graphs of graphs I

The following is from [LM09], which used an argument from an old (unpublished) version of this paper [Lou06]. A *graph of graphs* is a graph of spaces such that all vertex and edge spaces are finite combinatorial graphs and all edge maps are embeddings. A graph is 2-covered by a family of subgraphs if every edge is contained in exactly two elements of the cover, and a graph of graphs X is 2-covered if each vertex space is 2-covered by incident edge spaces. A graph of graphs X has an *underlying graph* $\Gamma_U(X)$, and corresponds to a map from $\Gamma_U(X)$ to the category of combinatorial graphs and injective maps. A *2-covered graph of graphs with boundary* is a graph of graphs such that non-valence-one vertex spaces are 2-covered by incident edge spaces, and valence one vertex spaces are isomorphic to their incident edge space.

Let $G = \Delta(F_i, Z_j)$ be a graph of free groups F_i over nontrivial subgroups Z_j . If $\phi: G \rightarrow \mathbb{F}$ is a homomorphism embedding each $F_i < G$, then there is a graph of spaces with fundamental group G and a natural map to a graph representing ϕ . For each i , choose an immersion $\varphi_i: \Gamma_i \rightarrow R$ where R is the rose with n petals, and fundamental group \mathbb{F} and $\pi_1(\Gamma_i) = F_i$, and Γ_i a core graph. For each j , choose an immersion $\varphi_j: \Delta_j \rightarrow R_n$ representing the image of Z_j . If $Z_j \hookrightarrow F_i$ then φ_j lifts to an immersion $\varphi_{i,j}: \Delta_j \rightarrow \Gamma_i$. Use the $\varphi_{i,j}$ as attaching maps of annuli, one for each edge of Δ , to define a graph of spaces X . The homomorphism ϕ induces a map $\varphi: X \rightarrow R$. Restricted to a product $\Delta_j \times I$, the map is projection to the first factor, followed by the immersion φ_j . Let b be the basepoint of R .

Let $\{V_p\}$ be the collection of connected components of $\varphi^{-1}(b)$, let edge graphs be connected components of preimages of midpoints of edges of R and regard X as a graph of graphs by regarding the V_p as vertex graphs.

The homomorphism $F_i \hookrightarrow G$ is represented by the inclusion $\Gamma_i \rightarrow X$, and the map $X \rightarrow R$ therefore represents the map $G \rightarrow \mathbb{F}$. Given $G \rightarrow \mathbb{F}$ as above, we sometimes write $X(G, \phi)$ to denote X . Since all vertical edges come from annuli X is clearly 2-covered.

Theorem 2.1. *Let $\pi: G \left[\sqrt[\iota]{\delta_j} \right] \rightarrow H$ be as in Theorem 1.2, and suppose further that G and H are free and of the same rank. Then the edge spaces of $X(G \left[\sqrt[\iota]{\delta_j} \right], \pi)$ are forests.*

The *horizontal* subgraph $\Gamma(X)$ of a graph of graphs X is the graph obtained by restricting to the zero skeleta of the vertex and edge spaces. Let $\Gamma_\infty(X)$ be the subset of $\Gamma(X)$ consisting of the connected components not homeomorphic to S^1 . For $X(G, \phi)$ as above, $\Gamma_\infty(X) = \cup_i \Gamma_i \subset X$. We assume throughout that if X is a graph of graphs without boundary, and e is an edge of a vertex space, then e is covered by at least two edges from incident edge spaces. This is equivalent to saying that the graph obtained by restricting to midpoints of edges of vertex and edge spaces is a core graph. An edge of a vertex or edge space is *vertical*.

2.3 Blowing up

Let X be a graph of graphs. If e is an embedded edge in the underlying graph of X then X/e , the *blow-down* of X , is the graph of graphs obtained by collapsing the edge space $E \times I$ to E . The quotient map $X \rightarrow X/e$ is a homotopy equivalence. The inverse of blowing down is *blowing up*.

Theorem 2.2 ([LM09, Theorem 2.4]). *Given a graph of graphs X there is a homotopy equivalent graph of graphs Y , obtained by repeatedly blowing up, such that each edge space of X is homeomorphic to a disjoint union of edge space of Y , vertex and edge spaces of Y are connected, and each vertex space of Y has exactly three incident edge spaces.*

Lemma 2.3 (Euler Characteristic Lemma). *Let X be a graph of graphs with connected vertex and edge spaces. Then*

$$\chi(\Gamma(X)) \leq \chi(\Gamma_U(X))$$

Suppose $\chi(\Gamma(X)) = \chi(\Gamma_U(X))$ and every vertex space has three incident edge spaces. If E_1, E_2 and E_3 are incident to a vertex space V with no monogons then $E_1 \cap E_2 \cap E_3$ is a single point and V is 2-covered.

Proof. Identical to the proof of Theorem 1.1 of [LM09]. Blow X up to a homotopy equivalent space Y with only valence three vertex spaces. Blowing

up and down don't change the Euler characteristic of horizontal or underlying graphs. If V is a vertex space with three incident edges, connectivity implies that there is a vertex of V in the image of all three incident edge spaces. This vertex corresponds to a valence three vertex of $\Gamma(X)$, therefore the map from valence three vertices of $\Gamma(X)$ to vertex spaces of X is surjective. Then by Gauss-Bonnet $\chi(\Gamma(X)) \leq \chi(\Gamma_U(X))$.

If $\chi(\Gamma(X)) = \chi(\Gamma_U(X))$ the map from valence three vertices of the horizontal graph to the set of valence three vertices of the underlying graph is bijective. If an edge of a vertex space is covered three times by incident edge spaces then the endpoints are covered three times by incident edge spaces. Since the valence three vertices of the underlying graph and the horizontal graph are in bijective correspondence, the endpoints of the offending edge must be the same, hence V contains a monogon. \square

Graphs of spaces whose horizontal subgraphs have the same Euler characteristic as their underlying graphs are particularly important. In this case vertex spaces have a simple local structure.

Lemma 2.4. *A graph V , 2-covered by connected subgraphs $E_i \subset V$, $i = 1, 2, 3$, such that $\cap \text{Im}(E_i)$ is a single vertex w , has one of the following forms:*

- E_i are all points.
- E_1 is a point, and $E_{2|3} \cong V$.
- $V = \text{Im}(E_1) \vee_w \text{Im}(E_2)$, $E_3 \cong V$.
- V is the union of three subgraphs $V_{1|2|3}$ which meet at a single vertex $w \in V \cong \bigvee_w V_i$. $E_i \cong V_{i+1} \vee_w V_{i+2}$.

If V is as above and $W \subset V$ we say that W is *cut by w* if W is not contained in some V_i . If $W_1, W_2 \subset V$ and neither W_1 nor W_2 is cut by w then we say that w separates W_1 and W_2 if, without loss of generality, $W_1 \subset V_1$, $W_1 \not\subset V_2$, and $W_2 \subset V_2$, $W_2 \not\subset V_1$.

Proof of Lemma 2.4. Let \mathcal{P} be the set of edge-paths starting at w and terminating if they meet w again. Let $\{E_i\}_{i=0,1,2}$ be the edge graphs incident to V . We divide \mathcal{P} into three subclasses $\mathcal{P}_{j,k}$, $j \neq k$. A path p lies in $\mathcal{P}_{j,k}$ if the image of p is contained in $\text{Im}(E_j) \cap \text{Im}(E_k)$. Let $V_i = \bigcup_{p \in \mathcal{P}_{i+1,i+2}} \text{Im}(p)$. At most one of V_i can be a point. Every point/edge of V lies in one of V_i , which all meet at w , the sole point of triple intersection. An incident edge E_i is then isomorphic to $V_{i+1} \cup V_{i+2}$. \square

2.4 Cylinders

A morphism of graphs of graphs is a morphism of graphs of spaces immersing vertex and edge spaces, respecting the product structure of edge spaces, in vertex and edge spaces, respectively. A *cylinder* in a graph of spaces X is a morphism $C \rightarrow X$, where C is a graph of graphs with all edge maps isomorphisms of connected graphs and with underlying graph a circle. Furthermore, the induced map $\Gamma_U(C) \rightarrow \Gamma_U(X)$ should be a cyclically reduced path. Cylinders are partially ordered by the relation “factors through”, i.e., $C \geq C'$ if $C' \rightarrow C \rightarrow X$.

Let $C \rightarrow X$ be a cylinder, and let F be a vertex space of C . If the map $F \rightarrow X$ is not an immersion, there are oriented edges h and k with the same initial endpoint with the same image in a vertex graph of X . Let $E \rightarrow F$ be an incident edge map. The edges e and f are the images of edges h' and k' in E . Since edge maps of C are all isomorphisms of graphs there is a rotation map $C \rightarrow C$ given by translating each vertex and edge space one unit: If edges are ordered coherently and (ι_i) and (τ_i) are the edge maps then the rotation map is given by $\tau_i \iota_i^{-1}$ on vertex spaces and by $\iota_{i+1}^{-1} \tau_i$ on edge spaces. If (a, b) is an element of the orbit of $(h^{(i)}, k^{(i)})$ then a and b are folded together by the associated map $C \rightarrow X$. Fold all pairs in the orbits of $(h^{(i)}, k^{(i)})$ to obtain a new cylinder $C' \rightarrow X$. The map $C \rightarrow X$ factors through the natural map $C' \rightarrow X$. Since folding is equivariant with respect to rotation $C' \rightarrow X$ is a cylinder.

Let $f_i: C_i \rightarrow X$ be a collection of cylinders. As in the case of a graph, we define the pullback of $\{C_i\}$ to be the graph of graphs

$$P = \{(x_1, \dots, x_n) \in C_1 \times \dots \times C_n \mid f_i(x_i) = f_j(x_j)\}$$

There are induced maps $g_i: P \rightarrow C_i$. If $C_1, C_2 \rightarrow X$ are cylinders and their pullback contains a cylinder then either, without loss, $C_1 \geq C_2$ or neither is maximal. Suppose Z is a cylindrical component of the pullback. Then $C' = (C_1 \cup C_2 \cup Z)/(g_1(z) \sim g_2(z))$ is a cylinder, and has an induced map $C' \rightarrow X$. Now fold to obtain a larger cylinder $C'' \rightarrow X$. If $C \rightarrow X$ is a cylinder, consider the pullback of the collection $C, C \rightarrow X$: if it contains a cylinder other than the diagonal then C is not maximal.

Let X be a graph of graphs, let I be a subdivided interval, and let $p: I \rightarrow X$ be a reduced edge path in $\Gamma(X)$. Let V be a vertex space of X , and say that $\eta: F \rightarrow V \subset X$ pushes along p if $p(0) = \eta(v)$ for some vertex v of F and the map $(I \cup F)/(0 \sim v) \rightarrow X$ extends to the product $I \times F$ and the reduced edge paths obtained by restricting to vertices of F are all identical.

Lemma 2.5. *Suppose $\chi(\Gamma(X)) = \chi(\Gamma_U(X))$, $\eta: F \looparrowright X$ pushes along p and p' , the first edges of the projections of p and p' to $\Gamma_U(X)$ agree, and that F has more than one point. Then, without loss of generality, the projection of p to $\Gamma_U(X)$ is a sub-path of the projection of p' to $\Gamma_U(X)$.*

Proof. Since F has more than one point, images of vertex spaces of $I \times F$ are uncut by valence three vertices of $\Gamma(X)$. We may assume by removing prefixes that the projections of p and p' diverge after one step, and that they are both of length two, but this implies that the image of F in the vertex space where the two paths diverge is cut by a valence three vertex of $\Gamma(X)$, contradicting the fact that F pushes past that vertex space. \square

Lemma 2.6. *Suppose $\chi(\Gamma(X)) = \chi(\Gamma_U(X))$. Suppose $\eta: F \looparrowright X$ pushes along a path p , and suppose $\eta(x) = \eta(y)$ for some $x \neq y$. Let $F' \looparrowright X$ be the immersed graph obtained by identifying x and y and folding. Then F' pushes along p .*

Proof. The two element set $\{x, y\}$ pushes along p , and $F/x \sim y$ therefore pushes along p . Furthermore, any two edges of $F/(x \sim y)$ adjacent to $[x]$ push along p , and since X is 2-covered, they push in a unique way, hence if e and f are identified by folding, then $(F/x \sim y)/(e \sim f)$ pushes along p . The lemma follows by induction on the number of edges. \square

Lemma 2.7 (Vertex spaces embed). *Let X be a 2-covered graph of spaces and let $C \rightarrow X$ be a cylinder. If E is an edge space of C and the induced map $E \rightarrow X$ is not injective then C is not maximal.*

Proof. Identify points having the same images and fold. \square

Lemma 2.8 (Vertex spaces do not overlap much). *Let $C, C' \rightarrow X$ be two cylinders such that all maps of vertex spaces are embeddings, and suppose two vertex spaces E and E' of C and C' , respectively, intersect in more than one point. Then either (without loss of generality), $C \geq C'$ or neither is maximal.*

Proof. The multi-element set $E \cap E' \subset X$ pushes along arbitrarily long reduced edge paths, but the pullback of C and C' is finite, therefore there is a cylindrical component of the pullback. \square

Existence of maximal cylinders is guaranteed by the previous corollaries. Let Y be obtained from X by blowing up. If C is a maximal cylinder in X then there is a maximal cylinder D in Y such that C is a blow-down of D and the two maps $D \rightarrow Y \rightarrow X$ and $D \rightarrow C \rightarrow X$ agree. In particular, the collection of types of cylinder cross-sections of X and Y are the same.

Lemma 2.9 (No cycles). *Let V be a vertex space of X , and suppose that C_0, \dots, C_{n-1} are distinct maximal cylinders, E_0, \dots, E_{n-1} vertex spaces of C_i , such that $C_i \rightarrow X$ embeds E_i in V . If $E_{(i \bmod n)} \cap E_{i-1} \neq \emptyset$ then $E_i \cap E_j \neq \emptyset$ for some $|i - j| > 1$. Intersections of images of edge graphs of maximal cylinders are cut-points in the associated edge spaces of X .*

Proof. Suppose not. Without loss of generality there are paths $p_{i,i+1}$, $i = 0..n-2$ such that $E_i \cap E_{i+1}$ pushes along $p_{i,i+1}$, the projection p of $p_{i,i+1}$ to $\Gamma_U(X)$ agree, and $E_0 \cap E_{n-1}$ doesn't push along any path projecting to p . Choose a longest path p' projecting to a sub-path of p that $E_0 \cap E_{n-1}$ pushes along. Then the images of E_0 and E_{n-1} are separated by the valence three vertex of the vertex space corresponding to the end of p' , but the relation 'not separated by' is transitive. Thus the sets $E_i \cap E_{i+1}$ push along arbitrarily long paths, and the pullback of C_i and C_{i+1} must contain a cylinder. \square




Figure 2: Edge spaces are trees of cylinder cross-sections.

Proof of Theorem 2.1. Consider the graph of graphs $X = X(G[\sqrt[l_j]{\delta_j}], \phi)$. Blow X up to a graph of graphs Y with only valence three vertices. Each edge space of X is a disjoint union of edge spaces of Y . We show that the edge spaces of Y are trees. Assuming $\text{rk}(G) = \text{rk}(H)$, we have $\chi(\Gamma(Y)) = \chi(\Gamma_U(Y))$. Let $M_j(\delta_j, l_j)$ be the subdivided mapping cylinder corresponding to δ_j , now regarded as an element of $\Gamma_\infty(Y)$. Then Y is obtained by gluing the boundaries of the mapping cylinders M_j to $\Gamma_\infty(Y)$ along reduced edge paths representing the δ_j . The map $M_j \rightarrow Y$ is a cylinder, and factors through a maximal cylinder C . Each δ_j is indivisible, therefore edge graphs of M_j embed in Y , hence edge graphs of M_j embed in C , and since no two M_j are conjugate or overlap in a vertical edge, C is the union of some subcollection of the M_j . Since the δ_j are nonconjugate, $C = M_j$ for some j . By Lemma 2.9 intersections of images of edge graphs of the M_j are cut points of edge spaces of Y . If e is an (vertical) edge of an edge space of Y then e is contained in the image of some cross section of some M_j , and

therefore each nontrivial edge space of Y is a union of trees along cut-points, hence is a tree. \square

Proof of Theorem 1.2 when G and H are free. By Theorem 2.1 edge spaces of X are forests, and since $l_j > 0$ there is some element δ_j which crosses an edge f of $\Gamma_\infty(X)$ only once, and no other $\delta_{j'}$ crosses f . Thus G can be written as $F * \langle \delta_j \rangle$ with all $\delta_{j'}$, $j' \neq j$, conjugate into F . Induct. \square

Proof of Theorem 1.3. The proof is exactly the same as the previous proof. Indivisibility of t in at least one factor implies that edge spaces of $X(G)$ are forests, and a valence one vertex of a forest corresponds to an edge that an immersion representing t in one of A or B crosses exactly once, hence t is primitive in at least one factor. \square

Proof of Theorem 1.4. The hypotheses are again designed to imply that edge spaces of $X(G, \phi)$ are forests. \square

3 Groups

3.1 Immersions II

Immersions of (core) graphs efficiently represent inclusions of free groups. We mimic this and represent injections

$$G = G_1 * \cdots * G_{p_G} * \mathbb{F}_{q_G} \hookrightarrow H = H_1 * \cdots * H_{p_H} * \mathbb{F}_{q_H}$$

with immersions of graphlike complexes.

Definition 3.1 (Relative Graph). A *relative graph* is a CW-complex Γ equipped with a pair of collections of disjoint connected subcomplexes $\text{Ell}(\Gamma) = \{Y_1, \dots, Y_p\} \subset \text{Verts}(\Gamma)$ such that cells not contained in any $Y \in \text{Ell}(\Gamma)$ are at most one-dimensional.

The *vertices* of a relative graph are the elements of $\text{Verts}(\Gamma)$, along with 0-cells not contained in any element of $\text{Verts}(\Gamma)$. The *edges* of a relative graph are the one-cells not contained in any element of $\text{Ell}(\Gamma)$. The valence of a vertex v is the number of edges with endpoints in v . A vertex is a *tree* if it has trivial fundamental group, and is *empty* if it is just a 0-cell. A *relative graph with boundary* is a relative graph with some valence one empty vertices marked as elements of $\partial\Gamma$. A relative graph Γ is reduced if all valence one empty vertices are in $\partial\Gamma$.

It will be convenient to also refer to the subcomplex $Y_1 \cup \dots \cup Y_p$ as $\text{Ell}(\Gamma)$. Denote by $\Gamma/\text{Ell}(\Gamma)$ the (ordinary) graph obtained by collapsing all elements of $\text{Ell}(\Gamma)$ to points. Let $\text{Ell}_T(\Gamma)$ be the set of elements of $\text{Ell}(\Gamma)$ with trivial fundamental group.

Definition 3.2. The complexity of a relative graph is the pair

$$\chi_{\text{fd}}(\Gamma) := (-\chi(\Gamma/\text{Ell}(\Gamma)), |\text{Ell}(\Gamma)|)$$

Pick a (empty or nonempty) vertex v . Take a copy of v , and for each edge e of Γ with terminal endpoint in v attach a copy of e at the corresponding point to obtain the *star* of v , $\text{St}(v)$. There is a natural map $\text{St}(v) \rightarrow \Gamma$.

The *inessential* vertices of a relative graph (with boundary) Γ are the empty valence-two vertices. The *essential* vertices are the non-boundary, non-inessential vertices. A relative graph Γ is *admissible* if stars of essential vertices embed and cover Γ . If S is a star of an essential vertex then we give S the structure of a relative graph with boundary by setting $\partial S = (\overline{\Gamma \setminus S}) \cap S$. An admissible relative graph is recovered by identifying stars of essential vertices in pairs. Let $\mathcal{N}(\Gamma)$ be the collection of all stars of essential vertices.

Definition 3.3. Let Γ and Γ' be relative graphs. A continuous map $\varphi: \Gamma \rightarrow \Gamma'$ is a morphism of relative graphs if elements of $\text{Ell}(\Gamma)$ are mapped to elements of $\text{Ell}(\Gamma')$, the preimage of the interior of an edge of Γ' is the union of interiors of edges from Γ , and the restriction of φ to the interior of such edge of Γ is a homeomorphism onto the corresponding interior in Γ' .

If v is a tree vertex of Γ and $\Gamma \rightarrow \Gamma'$ is a morphism of relative graphs, let b_1, \dots, b_n be the 0-cells of Γ contained in v which intersect an edge of Γ . Let v' be a tree with n boundary components b'_1, \dots, b'_n , and let Γ'' be the relative graph obtained by replacing v by v' , gluing the edges of Γ attached to v at b_i to b'_i . There are maps $\Gamma'' \rightarrow \Gamma \rightarrow \Gamma'$ such that $\Gamma'' \rightarrow \Gamma'$ is the identity away from v and v' , and an isomorphism of fundamental groups. We make this adjustment without comment.

Definition 3.4 (Immersion (binding ties [Sta65])). Let $\varphi: \Gamma' \rightarrow \Gamma$ be a combinatorial map of relative graphs, and suppose that Γ is admissible. We say that φ is an *immersion* if, for all vertices of Γ and connected components w of $\varphi^{-1}(v)$,

$$\pi_1(\varphi_w): \pi_1(\text{St}(w), \partial \text{St}(w)) \rightarrow \pi_1(\text{St}(v), \partial \text{St}(v))$$

is injective.




Figure 3: Homotopy between reduced edge paths is a concatenation of rectangles.

Let Γ be a relative graph. A *basic rectangle* in Γ is a map $[0, 1] \times [0, 1] \rightarrow \Gamma$ such that $\{0\} \times [0, 1]$ and $\{1\} \times [0, 1]$ are mapped to 0-cells meeting edges in Γ , and with image either completely contained in an element of $\text{Ell}(\Gamma)$ or such that the sets $(0, 1) \times \{y\}$ are mapped homeomorphically to interiors of edges of Γ . A *rectangle* in Γ is either a rectangle of the second type or a map of a square $[0, 3] \times [0, 1] \rightarrow \Gamma$ such that $[0, 1] \times [0, 1] \rightarrow \Gamma$ and $[2, 3] \times [0, 1] \rightarrow \Gamma$ are basic rectangles of the second type and $[1, 2] \times [0, 1] \rightarrow \Gamma$ is a basic rectangle of the first type. If R is a rectangle define $\partial^L(R) = \{0\} \times [0, 1]$, $\partial^R(R) = \{1\} \times [0, 1]$, and $\partial^+(R) = [0, 1] \times \{1\}$ and $\partial^-(R) = [0, 1] \times \{0\}$.

An *edge path* in a relative graph is a morphism of a subdivided interval. A *reduced edge path* is an immersion. An edge path is homotopic, relative to its endpoints, to a reduced edge path, and any two reduced edge paths homotopic relative to endpoints are homotopic through reduced edge paths, i.e., they differ by reparametrization of the interval and a homotopy supported on subsegments mapping to elements of $\text{Ell}(\Gamma)$. See Figure 3. A homotopy between two reduced edge paths is a concatenation of basic rectangles.

The following lemma follows from the innermost disc argument.

Lemma 3.5. *Let Γ be a graph, let $p_0, p_1: S^1 \rightarrow \Gamma$ be two homotopic reduced edge paths, and let $P: A \rightarrow \Gamma$ be a map of an annulus $A = S^1 \times I$ such that $P|_{S^1 \times \{i\}} = p_i$. Then there is a map $Q: A \rightarrow \Gamma$ which restricts to p_0 and p_1 on the boundary and is a cycle of basic rectangles.*

A map of an annulus as in the lemma is *admissible*. An admissible annulus is as a homotopy between two cyclically reduced paths through




Figure 4: Folding.

cyclically reduced paths, similar to the case of a homotopy between reduced edge paths through reduced edge paths.

Definition 3.6 (Folding). Folding: Let Γ be a relative graph, let v be a vertex of Γ , and let e and f be *distinct* edges with terminal endpoints in v . Let D be a square, and attach three consecutive sides of D to Γ by a path $ep\bar{f}$, where p is a path in v . (Recall that v may have nontrivial fundamental group.) Collapse D by collapsing the face attached to f to the remaining three sides, and if the fourth side of D connects two 0-cells of Γ crush the fourth side to a point to obtain Γ' . The vertices of Γ' are obtained from those of Γ by taking connected components of the image of $\text{Ell}(\Gamma)$ and the fourth side of D .

Attach 2-cells: Let Γ be a relative graph. If $Y \in \text{Ell}(\Gamma)$ attach 2-cells to Y .

Collapsing: Replace an element Y of $\text{Ell}(\Gamma)$ by a subcomplex of Γ containing Y . Collapsing to a point: Replace $Y \in \text{Ell}(\Gamma)$ by a point.

The first kind of fold is illustrated in Figure 4. If v is an ordinary vertex then the path p is trivial and a fold is just a Stallings fold.

Lemma 3.7 ([Sta83]). *If $\varphi: \Gamma \rightarrow \Gamma'$ is combinatorial, then there is a relative graph $\bar{\Gamma}$, $\chi_{\text{fd}}(\bar{\Gamma}) \leq \chi_{\text{fd}}(\Gamma)$, a π_1 -onto map $F: \Gamma \rightarrow \bar{\Gamma}$, and an immersion $\bar{\varphi}: \bar{\Gamma} \rightarrow \Gamma'$ such that $\bar{\varphi} \circ F$ is homotopic to φ , and F is a composition of folds.*

Immersions are injective on fundamental group.

The proof of Lemma 3.7 is an easy variation on Stallings/Dunwoody folding sequences.

Proof. If φ maps a nonempty vertex to a point, collapse the vertex to a point and remove it from the collection $\text{Ell}(\Gamma)$ (if it is in $\text{Ell}(\Gamma)$). The complexity strictly decreases in the second case.

Otherwise, if φ isn't an immersion there is a vertex v' of Γ' and a connected component v of $\varphi^{-1}(v')$ such that the induced map of fundamental groups of stars is not injective. Either the map extends to $\Gamma \cup D$, as in Definition 3.6 and we fold, and if not, attach enough two-cells to $\text{Ell}(\Gamma)$ so the restriction is injective on (ordinary) fundamental group. If the fourth edge connects distinct nonempty vertices of Γ then the complexity decreases, and if the fourth edge connects a vertex to itself, the complexity also decreases.

The last claim follows from a standard innermost disc argument. \square

One important property of immersions of relative graphs is that if p is a reduced edge path then the composition of p and an immersion is also a reduced edge path. More generally, compositions of immersions are immersions.

If $\varphi: \Gamma \rightarrow \Gamma'$ is an immersion of relative graphs and Γ' is admissible, $S \subset \Gamma'$ a star of an essential vertex, then the connected components of $\varphi^{-1}(S)$ are (embedded!) stars of connected subcomplexes of Γ . Without loss of generality, we assume that if such a subcomplex contains an element of $\text{Ell}(\Gamma)$ then it is in $\text{Ell}(\Gamma)$. This may decrease the cardinality of $\text{Ell}(\Gamma)$.

3.2 Graphs of graphs II

In this subsection we consider the situation of Theorem 1.2. The objective of this section is to prove the following theorem, which we reduce to an analysis of graphs arising by adjoining roots to free groups.

For the remainder of this section, fix an inclusion $\eta: G \hookrightarrow H$ of finitely generated groups which lifts to an epimorphism $\pi: G' \twoheadrightarrow H$. Before we begin we replace H by a group H' as follows. Let H' be a group of maximal χ_{fd} such that $G' \twoheadrightarrow H$ factors through $G' \twoheadrightarrow H'$. By Lemma 3.7:

Lemma 3.8. *Let $H' \twoheadrightarrow H$ be as above. Then*

$$\chi_{\text{fd}}(H') \geq \chi_{\text{fd}}(H)$$

If a freely indecomposable free factor of H' has trivial image in H then the inequality is strict.

We represent η as an immersion $\varphi: \Gamma_G \rightarrow \Gamma_H$ and construct a space X , similar to a graph of graphs, with fundamental group $G \left[\begin{smallmatrix} k_i/\gamma_i, \\ l_i/\delta_j \end{smallmatrix} \right]$. This space has a nice map to Γ_H representing π . Furthermore, by resolving, as

in the case of a free group, X may be given a graph of spaces structure “transverse” to the old one. After adjusting X and analyzing the structure of vertex spaces we reduce the proof of Theorem 1.2 to Theorem 2.1.

Choose an admissible relative graph Γ_H with fundamental group H , and whose components $Y_1, \dots, Y_p \in \text{Ell}(\Gamma_H)$ have fundamental groups $\pi_1(Y_i) \cong H_i$. Choose a relative graph Γ_G with fundamental group G and an immersion $\varphi: \Gamma_G \looparrowright \Gamma_H$ provided by Lemma 3.7 representing η . Choose, for each γ_i, δ_j , immersions $\gamma_i: S^1 \looparrowright \Gamma_G, \delta_j: S^1 \looparrowright \Gamma_G$, representing the conjugacy classes $[\gamma_i], [\delta_j]$. Each immersion $\gamma_i: S^1 \rightarrow \Gamma_G$ has image in the connected component of $\text{Ell}(\Gamma_G)$ representing G_i .

Let M_j be the mapping cylinder of the l_j fold cover $S_j^1 \rightarrow S^1$, and let r_j be the core curve (the codomain S^1) of M_j . Now glue the M_j along S_j^1 to Γ_G using the immersions δ_j as attaching maps to form a space X . For each r_j , choose an immersion $\sqrt[l_j]{\delta_j}: r_j \rightarrow \Gamma_H$ representing the conjugacy class of $\sqrt[l_j]{\delta_j} \in H$. The map $\varphi: \Gamma_G \looparrowright \Gamma_H$ lifts to a continuous map $\pi: X \rightarrow \Gamma_H$ agreeing with the immersions r_j and φ . Each mapping cylinder M_j is a quotient space of an annulus A_j , and by Lemma 3.5 we may assume that π induces admissible maps of annuli.

The space X is a quotient of the disjoint union of annuli, Γ_G , and the core curves r_j of the mapping cylinders M_j . Set $\Gamma(X) = \Gamma_G \sqcup \bigsqcup r_j$; $\Gamma_G(X)$ plays the role that $\Gamma_\infty(X)$ does in the free case.

As with graphs of free groups, we give X a graph of spaces structure transverse to the graph of spaces decomposition $\Gamma_G \sqcup \bigsqcup M_j / \sim$. Let \mathcal{Z} be the collection of connected components of $\pi^{-1}(N)$, $N \in \mathcal{N}(\Gamma_H)$, and let \mathcal{B} be the collection of connected components of the preimages of valence two vertices. Since each mapping cylinder induces an admissible map of the associated annulus the elements of \mathcal{B} are graphs contained in X . For each $B \in \mathcal{B}$ there are two embeddings $B \hookrightarrow \mathcal{Z}$. Since each γ_i is conjugate into some G_j the associated mapping cylinders M_i are completely contained in the Z . The boundary of $Z \in \mathcal{Z}$ is the set $\mathcal{B} \cap Z$, and coincides with the set of points of Z mapping to valence two 0-cells of Γ_H .

Each mapping cylinder is either completely contained in some $Z \in \mathcal{Z}$ or has nontrivial intersection with \mathcal{B} . If M has nontrivial intersection with \mathcal{B} then $r \cap Z$ (recall that r is the core curve of M) is a collection of closed intervals, and the preimage \mathcal{B} slices an annulus $A \in \mathcal{A}$ into rectangles.

The connected components of the intersections of the relative graph $\Gamma(X)$ and elements of \mathcal{Z} are relative graphs S with distinguished valence one vertices $S \cap \mathcal{B}$. Let $\mathcal{S}(Z)$ be the collection of connected components of $\Gamma(X) \cap Z$. If $S \in \mathcal{S}$ set $\partial S = S \cap \mathcal{B}$. For a fixed $Z \in \mathcal{Z}$ let $\mathcal{R}(Z)$ be the

subcollection of all rectangles contained in Z .

The boundary of each rectangle R is composed of two types of arcs, $\partial^\pm R = R \cap \Gamma(X)$ and $\partial^{L|R} R = R \cap \mathcal{B}$. The former are *horizontal boundary arcs* and the latter are *vertical boundary arcs*.

For each rectangle R let $\varphi_R^\pm: \partial^\pm R \rightarrow S(R, \pm)$

the attaching maps for the horizontal/vertical boundary arc of R . Then Z is the quotient space

$$\mathcal{S}(Z) \sqcup \mathcal{R}(Z) / (x \sim \varphi_R^+(x), y \sim \varphi_R^-(y))$$

The boundary of Z is the union of vertical boundary arcs of rectangles contained in Z .

We reconstruct Z by gluing the rectangles $\mathcal{R}(Z)$ to $\mathcal{S}(Z)$ via the attaching maps $\varphi_R^\pm: (\partial^\pm R, \partial\partial^\pm R) \rightarrow \sqcup(S, \partial S) \in \mathcal{S}(Z)$.




Figure 5: Crushing collars of boundary components.

The boundary of Z is the union of vertical boundary arcs of the rectangles comprising it, along with all valence one vertices of $\partial\mathcal{S}(Z)$ not contained in some vertical boundary arc of a rectangle. By construction, Z is a connected component of $\varphi^{-1}(N)$ for some $N \in \mathcal{N}(\Gamma_H)$. Let $\{b_1, \dots, b_k\}$ be the valence one vertices comprising ∂N , with incident edges $e_i \subset N$ so that $\tau(e_i) = b_i$. If $\psi: Z \rightarrow N$ is the restriction of π then $\psi^{-1}(e_i)$ is a collar neighborhood of $\psi^{-1}(b_i) \in \mathcal{B}$, a boundary component of Z . The restriction ψ factors through the map which projects each collar onto the I factor. Call the resulting quotient space \bar{Z} .

The map $\pi: X \rightarrow \Gamma_H$ factors through the graph of spaces Γ_K obtained by assembling the collection $\{\bar{Z} \mid Z \in \mathcal{Z}\}$ along images of boundary components: if $B \subset Z_1, Z_2$, then identify the images $\bar{B} \in \bar{Z}_1$ and $\bar{B} \in \bar{Z}_2$.

Lemma 3.9. *The \bar{Z} have freely indecomposable (or trivial!) fundamental groups.*

Proof. Suppose not. Then $\pi_1(X) \twoheadrightarrow H$ factors through a group with strictly higher Scott complexity. If some \bar{Z} had freely decomposable fundamental group then $\chi_{\text{fd}}(K) > \chi_{\text{fd}}(H)$. \square

If $q_H = q_G$, then no $S \in \mathcal{S}(Z)$ has fundamental group with nontrivial free part. In particular, elements of $\mathcal{S}(Z)$ are all relative trees. Let $G = G(Z)$ be the graph with vertex set $\mathcal{S}(Z)$ and edge set $\mathcal{R}(Z)$. The endpoints of an edge R are the boundary components $\partial^\pm R$, and an endpoint $\partial^\pm R$ is attached to S if the image of φ^\pm is contained in S . Let T be a maximal tree in G and let Z_T be the space obtained by restricting to T .

Since no $S \in \mathcal{S}(Z)$ has nontrivial free part and G embeds in H , the components $Z \in \mathcal{Z}$ fall into three classes:

$$\mathcal{Z}_1 = \{Z \mid \pi_1(\bar{Z}) \text{ is trivial}\}.$$

$$\mathcal{Z}_2 = \{Z \mid \pi_1(\bar{Z}) \text{ is nontrivial and } Z \text{ contains no } Y \in \text{Ell}(\Gamma_G)\}$$

$$\mathcal{Z}_3 = \{Z \mid \pi_1(\bar{Z}) \text{ is nontrivial and } Z \text{ contains some } Y \in \text{Ell}(\Gamma_G)\}$$

Since $G \hookrightarrow H$ clearly if $Z \in \mathcal{Z}_1$ then Z contains no $Y \in \text{Ell}(\Gamma_G)$.

We reduce the computation $\chi_{\text{fd}}(H) \leq \chi_{\text{fd}}(G)$ to a version of the combinatorial Gauss-Bonnet Theorem.

Definition 3.10 (*q-excess/deficiency*). For each Z , let

$$\Delta_q^-(Z) = \frac{1}{2} (\#\partial Z_T - \#\partial \bar{Z})$$

and for $Z \in \mathcal{Z}_3$ let

$$\Delta_q^+(Z) = \frac{1}{2} b_1(\partial Z_T)$$

and for $Z \in \mathcal{Z}_1 \cup \mathcal{Z}_2$ set $\Delta_q^+(Z) = 0$.

If S is a relative graph with no loops (no contribution to q_G), set $\kappa(S, \partial S) = \frac{1}{2} \#\partial S - 1$,

Definition 3.11 (*p-excess/deficiency*). If $Z \in \mathcal{Z}_1$ then define $\Delta_p^+(Z) = \Delta_p^-(Z) = 0$. If $Z \in \mathcal{Z}_2$ then define $\Delta_p^+(Z) = 1$ and $\Delta_p^-(Z) = 0$. If $Z \in \mathcal{Z}_3$ and Y_0, \dots, Y_k are the components of $\text{Ell}(\Gamma_G)$ contained in Z , then define $\Delta_p^-(Z) = k$ and $\Delta_p^+(Z) = 0$.

Lemma 3.12.

$$\kappa(\bar{Z}, \partial \bar{Z}) - \sum_{S \in \mathcal{S}(Z)} \kappa(S, \partial S) = \Delta_q^+(Z) - \Delta_q^-(Z)$$

Proof. We need to show that

$$\frac{1}{2}\#\partial Z_T - 1 = \sum_{S \in \mathcal{S}(Z)} \left(\frac{1}{2}\#\partial S - 1 \right) + \Delta_q^+(Z)$$

Let $T \subset G(Z)$ be a maximal tree, and let S_1, \dots, S_k be an enumeration of $\mathcal{S}(Z)$ such that S_{i+1} is connected to $S_0 \cup R_0 \cup S_1 \cup R_1 \cdots S_i$ by an edge $R_i \subset T$. Assume that R_i is oriented so that $\partial^+ R_i$ is attached to S_{i+1} . Let Z_i be the union of S_1, \dots, S_i and R_1, \dots, R_{i-1} .

The boundary of $\partial^{\pm} R_i$ consists of two components. Suppose that for at least one of $+$ or $-$, the image $\partial\partial^{\pm} R_i$ is contained in two *distinct* boundary components of at least one of Z_i or S_{i+1} . If this is the case then

$$\begin{aligned} \kappa(Z_{i+1}, \partial Z_{i+1}) &= \frac{1}{2}\#\partial Z_{i+1} - 1 \\ &= \frac{1}{2}(\#\partial Z_i + \#\partial S_{i+1} - 2) - 1 \\ &= \kappa(Z_i, \partial Z_i) + \kappa(S_{i+1}, \partial S_{i+1}) \end{aligned}$$

If both $\partial\partial^+ R_i$ and $\partial\partial^- R_i$ have image in the same boundary component of S_{i+1} , Z_i , respectively, then Z_i contains at least $b_1(\partial Z_i) + 1$ elements of $\text{Ell}(\Gamma_G)$ and S_{i+1} must contain a new element of $\text{Ell}(\Gamma_G)$. We then have

$$\begin{aligned} \kappa(Z_{i+1}, \partial Z_{i+1}) &= \frac{1}{2}\#\partial Z_{i+1} - 1 \\ &= \frac{1}{2}(\#\partial Z_i + \#\partial S_{i+1} - 1) - 1 \\ &= \kappa(Z_i, \partial Z_i) + \kappa(S_{i+1}, \partial S_{i+1}) + \frac{1}{2} \end{aligned}$$

Each such rectangle makes a contribution of $+1$ to $b_1(\partial Z_T)$, $+1/2$ to Δ_q^+ , and only such rectangles make such contributions, thus

$$\kappa(Z_T, \partial Z_T) = \sum_{S \in \mathcal{S}(Z)} \kappa(S, \partial S) + \Delta_q^+(Z)$$

□

Lemma 3.13. *If $Z \in \mathcal{Z}_3$ then $\Delta_p^-(Z) \geq b_1(\partial Z_T) = 2\Delta_q^+(Z)$.*

Consider Figure 6. The tree in this example is a tripod, there is one boundary component of Z_T , and $\Delta_p^- = 2$. A similar example with only two components S would have $\Delta_p^- = 1$. We argue that each independent loop in ∂Z_T must contribute *at least* 1 to Δ_p^- .




Figure 6: Illustration for Lemma 3.13.

Proof of Lemma 3.13. Argue as in Lemma 3.12. If $\partial\partial^\pm R$ maps to a single boundary component of $S \in \mathcal{S}(Z)$ then, since $\partial^\pm R \rightarrow S$ is an immersion, S must have nontrivial fundamental group, and since the free part of S is trivial, it must contain some element Y of $\text{Ell}(\Gamma_G)$.

Choose the exhaustion of T so that S_0 has an incident edge R , $\partial^- R \rightarrow S_0$ such that $\partial\partial^- R$ maps to a single boundary vertex of S_0 . Let $R_{i_1}, \dots, R_{i_{b_1(\partial Z_T)}}$ be the rectangles such that $\partial\partial^+ R_{i_j}$ maps to a single boundary component of S_{i_j+1} . Then each S_{i_j+1} contains some element $Y_j \in \text{Ell}(\Gamma_G)$. Since the S_{i_j} are distinct, Z contains at least $b_1(\partial Z_T) + 1$ elements of $\text{Ell}(\Gamma_G)$, i.e., $\Delta_p^-(Z) \geq b_1(\partial Z_T)$. \square

Lemma 3.14. *If $Z \in \mathcal{Z}_2$ then $\Delta_q^-(Z) \geq \frac{1}{2}$. If equality holds then $\pi_1(\bar{Z}) \cong \mathbb{Z}_2$.*

Proof. Suppose that $\Delta_q^-(Z) > 0$. Clearly $\Delta_q^-(Z) \geq \frac{1}{2}$. Suppose $\Delta_q^-(Z)$ is $\frac{1}{2}$. Let Z_T be as before, and attach a rectangle R to Z_T such that $\#\partial Z_T \cup R = \#\partial Z_T - 1$. Let R_1, \dots, R_n, S^\pm be a path in T connecting the endpoints of R . Then R and the R_i must form a Mobius strip, as in Figure 7. The boundary is in bold. Adding rectangles while maintaining $\Delta_q^-(Z) = \frac{1}{2}$ does not change the fundamental group of \bar{Z} , which is clearly \mathbb{Z}_2 . \square

Lemma 3.15. $\chi_{\text{fd}}(H) \leq \chi_{\text{fd}}(G)$. *If equality holds then for all $Z \in \mathcal{Z}_2$, $\Delta_q^-(Z) = \frac{1}{2}$. In particular, every such \bar{Z} has fundamental group \mathbb{Z}_2 .*




Figure 7: Collapsing a collar neighborhood of a Mobius strip to build a projective plane.

If H has no \mathbb{Z}_2 free factors and equality holds then \mathcal{Z}_2 is empty, all members of \mathcal{Z}_3 contain exactly one element of $\text{Ell}(\Gamma_G)$, and $\Delta_{p|q}^\pm(Z) = 0$ for all Z .

Proof. By Gauss-Bonnet,

$$\sum_{Z \in \mathcal{Z}} \sum_{S \in \mathcal{S}(Z)} \kappa(S, \partial S) = q_G - 1$$

and, since the maps $\bar{Z} \rightarrow H$ don't factor through free products, again by Gauss-Bonnet,

$$\sum_{Z \in \mathcal{Z}} \kappa(\bar{Z}, \partial \bar{Z}) = q_H - 1$$

By homology, $q_H - 1 \leq q_G - 1$, and if $q_H - 1 = q_G - 1$, then by Lemma 3.12,

$$\sum_{Z \in \mathcal{Z}_3} \Delta_q^+(Z) = \sum_{Z \in \mathcal{Z}} \Delta_q^-(Z),$$

and we compute

$$\begin{aligned}
p_G &= |\mathcal{Z}_3| + \sum_{Z \in \mathcal{Z}_3} \Delta_p^-(Z) && \text{(Definition)} \\
&\geq |\mathcal{Z}_3| + 2 \sum_{Z \in \mathcal{Z}_3} \Delta_q^+(Z) && \text{(Lemma 3.13)} \\
&= |\mathcal{Z}_3| + 2 \sum_{Z \in \mathcal{Z}} \Delta_q^-(Z) && \text{(Lemma 3.12)} \\
&= |\mathcal{Z}_3| + 2 \sum_{Z \in \mathcal{Z}_2} \Delta_q^-(Z) && \text{(obvious)} \\
&\geq |\mathcal{Z}_3| + |\mathcal{Z}_2| && \text{(Lemma 3.14)} \\
&= p_H && \text{(Lemma 3.9)}
\end{aligned}$$

□

Each $Z \in \mathcal{Z}_3$ is a union of stars of vertices of $\Gamma(X)$, rectangles from mapping cylinders representing roots adjoined to elements δ_j , and mapping cylinders representing roots adjoined to elements γ_i which are conjugate into freely indecomposable free factors of G . Let $Y(Z)$ be the only element of $\text{Ell}(\Gamma_G)$ contained in Z , let $S_0(Z)$ be the element of $\mathcal{S}(Z)$ containing $Y(Z)$, and let $\mathcal{M}(Z)$ be the collection of mapping cylinders attached to $Y(Z)$. Let $\mathcal{R}(Z)$ again stand for the rectangles contained in Z . For each attaching map $p: \partial^\pm R \rightarrow (S, \partial S)$ we choose a fixed reduced edge path $\hat{p}: (I_p, \partial I_p \rightarrow (S, \partial S))$ such that if $[p] = [p'] \in \pi_1((S, \partial S))$ then there are maps $q: \partial^\pm R \rightarrow (S, \partial S)$ and $q': \partial^\pm R' \rightarrow (S, \partial S)$ such that $\hat{p}q = p$ and $\hat{p}q' = p'$.

Define an equivalence relation \sim on the collection of rectangles R by $R \sim R'$ if $[\partial^\pm R \rightarrow (S, \partial S)] = [\partial^\pm R' \rightarrow (S, \partial S)]$ and R and R' are not separated by a cutpoint in $G(Z)$.

The following is the analog of Lemma 2.4 for the vertex spaces $Z \in \mathcal{Z}_3$.

Lemma 3.16 (Structure theorem for vertex spaces.). *Let $Z \in \mathcal{Z}_3$ and suppose that*

- $S_i \in \mathcal{S}(Z)$ is a tree unless $i = 0$
- The fundamental group of S_0 embeds under the map $Z \rightarrow Z/\partial Z$

Then

- If $S \in \mathcal{S}$ is not a cutpoint in $G(Z)$ then there is only one homotopy class of attaching map associated to S .

- The \sim equivalence classes are exactly the maximal cut-point-free subgraphs of $G(Z)$.

Proof. Consider a chain of rectangles R_0, \dots, R_{n-1} forming an embedded cycle in $G(Z)$, and orient the edges of $G(Z)$ so that, for each i , $\partial^+ R_i$ and $\partial^- R_{i+1}$ are mapped to the same vertex space S_i . If $\partial^+ R_i$ and $\partial^- R_{i+1}$ are attached to the same S_{j_i} , $j_i \neq 0$, then the maps must agree up to reversal of orientation since $\Delta_q^-(Z) = 0$. Orient all rectangles so that $[\partial^+ R_i \rightarrow S_{j_i}] = [\partial^- R_{i+1} \rightarrow S_{j_i}]$ agree. Then the concatenation of rectangles, in $Z/\partial Z$ is a homotopy between $[\partial^+ R_0 \rightarrow S_{j_0}]$ and $[\partial^- R_{n-1} \rightarrow S_{j_0}]$, and since $\pi_1(S_0)$ embeds under $Z \rightarrow Z/\partial Z$, the attaching maps are equal.

If S is not a cutpoint then every pair of edges adjacent to S is contained in an embedded cycle in $G(Z)$. Any maximal cutpoint-free subgraph contains an embedded loop passing through any pair of edges, hence the second bullet holds as well. \square

Let $Z \in \mathcal{Z}_3$ be as above. Let B_i be the maximal cut-point-free subgraphs of $G(Z)$. For each class of attaching map $p: \partial^\pm R \rightarrow (S, \partial S)$ let I_p be an interval as before, and let Z_i be the graph of spaces obtained by replacing each $S \in B_i$ by I_p . Let p_1, \dots, p_k be the attaching maps for rectangles adjacent to $S_0 = \text{St}(Y(Z))$, where $Y(Z)$ is the element of $\text{Ell}(\Gamma_G)$ contained in S_0 . Let e_1, \dots, e_l be the oriented edges of X terminating in Y . Each path p_i is a concatenation of paths $e_{a(i)} r_i \bar{e}_{b(i)}$. Construct a new graph S'_0 by identifying the initial and terminal segments of I_{p_i} mapping to the same edges of $\text{St}(Y)$. We recover S_0 by attaching S'_0 to $Y(Z)$ by gluing the subgraph D_Z of S'_0 associated to the middle segments of the I_p along the paths r_i . There is a natural map $I_p \rightarrow S'_0$. Define a related space V_Z by taking $\mathcal{S}(V_Z) = \{S'_0, S_1, \dots, S_n\}$ as vertex spaces and attaching the rectangles from $\mathcal{R}(Z)$ to $\mathcal{S}(V_Z)$ with the induced attaching maps. Then Z is recovered by gluing the image of D_Z in V_Z to Y along the map $\psi_Z: D_Z \rightarrow Y(Z)$ induced by the r_i and reattaching the mapping cylinders in $\mathcal{M}(Z)$ to Y .

Each Z_i is homeomorphic to a product $B_i \times I$. Give $B_i \times I$ the foliation induced by projection to the I coordinate, and give each $S \in \mathcal{S}(V_Z)$ the foliation whose leaves are simply the points of S . Then define $F(Z)$ to be the foliation on V_Z induced by the foliations on $B_i \times I$ and S . Define $\Gamma_U(V_Z)$ to be the leaf space of $F(Z)$, and denote the quotient map by π_Z . Let $\partial\Gamma_U(V_Z)$ be the image of the boundary of Z . The following lemma is obvious from the construction.

If Γ is a graph with boundary $\partial\Gamma$, define

$$\kappa(\Gamma, \partial\Gamma) = -\chi(\Gamma) + \frac{1}{2}\#\partial\Gamma$$




Figure 8: V_Z and all the rest.

Lemma 3.17. *The following hold for $\Gamma_U(V_Z)$:*

- $\Gamma_U(V_Z)$ is a graph.
- $\kappa(\Gamma_U(V_Z), \partial\Gamma_U(V_Z)) = \sum_{S \in \mathcal{S}(V_Z)} \kappa(S, \partial S)$
- Each boundary component of $\Gamma_U(V_Z)$ is the image of exactly one boundary component of Z' .
- Point preimages under π_Z are connected and the natural maps $(S, \partial S) \rightarrow (\Gamma_U(V_Z), \partial\Gamma_U(V_Z))$ are immersions.

The following extension property also holds. Let $f: Z \rightarrow A$ be a continuous map to an aspherical space A , mapping each boundary component of Z to a point. Then there exists $f': \Gamma_U(V_Z) \rightarrow A$ such that $f' \circ \pi_Z$ and f are homotopic rel ∂Z . In particular, they induce the same map on $\pi_1(Z, \partial Z)$.

Let N_Z be the space obtained by attaching $\Gamma_U(V_Z)$ and all mapping cylinders in $\mathcal{M}(Z)$ to $Y(Z)$, and let \mathcal{N} be the collection of all such N_Z . The space X is the pushout of the diagram $\mathcal{B} \hookrightarrow \mathcal{Z}$. The map $\mathcal{B} \hookrightarrow \mathcal{Z}$ factors through the natural map $\mathcal{B} \hookrightarrow \mathcal{V}$, and the map $X \rightarrow \Gamma_H$ factors through the map crushing all elements of \mathcal{B} to points. By Lemma 3.17 $\bar{\mathcal{B}} \rightarrow \bar{\mathcal{Z}}$ factors through the natural map $\bar{\mathcal{B}} \rightarrow \mathcal{N}$, and by Van Kampen, $X \rightarrow \Gamma_H$ then factors, up to homotopy, through the pushout of $\bar{\mathcal{B}} \rightarrow \mathcal{N}$.

Define a new graph of graphs X' as the pushout of $\mathcal{B} \rightarrow \mathcal{V}$. The underlying graph of X' is obtained as the pushout $\bar{\mathcal{B}} \rightarrow \sqcup \Gamma_U(V_Z)$. Then $\Gamma(X')$ is the pushout $\mathcal{B}^0 \rightarrow \sqcup \mathcal{S}(V_Z)$.

It's clear that $\Gamma(X')$ and $\Gamma_U(X')$ are graphs, and, since point preimages of $V_Z \rightarrow \Gamma_U(V_Z)$ are connected, $\pi_1(X') \rightarrow \pi_1(\Gamma_U(X'))$ is onto. The attaching

maps $M \rightarrow \Gamma_G$ factor through the inclusion $\Gamma_\infty(X') \hookrightarrow X'$, and we have $X = \Gamma_\infty(X) \cup_{\delta_j} \{M_j\}$.

We recover X in the same way as we recover Z by gluing each $D_Z \subset \Gamma(V_Z)$ to $Y(Z)$ via the map ψ_Y . Since each $\Gamma(V_Z)$ immerses in Γ_Z under π_Z $\Gamma(X) \rightarrow \Gamma_U(X)$ is an immersion.

Lemma 3.18. $\chi(\Gamma(X')) = \chi(\Gamma_U(X'))$

Proof. The proof is the straightforward computation. By Lemma 3.17 and definition,

$$\begin{aligned} -\chi(\Gamma(X')) &= \sum_{Z \in \mathcal{Z}} \sum_{S \in \mathcal{S}(V_Z)} \kappa(S, \partial S) \\ &= \sum_{Z \in \mathcal{Z}} \kappa(\Gamma_U(V_Z), \partial \Gamma_U(V_Z)) \\ &= -\chi(\Gamma_U(X')) \end{aligned}$$

□

The images of the D_Z are disjoint under the map $X' \rightarrow \Gamma_U(X')$. We now prove Theorem 1.2.

Proof of Theorem 1.2. It suffices to show that G splits as $G' * \langle \delta_1^{d_1} \rangle * \dots * \langle \delta_m^{d_m} \rangle$. By 2.1 and connectivity of point preimages under $V_Z \rightarrow \Gamma_U(V_Z)$ edge spaces of X' are trees. Since the images of the D_Z are disjoint under the map $X' \rightarrow \Gamma_U(X')$ and the δ_j are not supported on D_Z , there is an edge of $\Gamma_\infty(X') \setminus \cup D_Z$ crossed by exactly one δ_j , and δ_j is therefore conjugate to a primitive element of G relative to G_i . Removing this edge from $\Gamma_\infty(X')$, along with the core curve of the mapping cylinder corresponding to δ_j and all rectangles adjacent to the core curve gives a graph of spaces of the same form. □

Remark 3.19. Simply knowing that some δ_j is a primitive element of $\pi_1(\Gamma(X))$ isn't sufficient to imply that $\langle \delta_j \rangle$ is a free factor of G , thus Theorem 1.2, even in the case $m = 1$, cannot be deduced from Baumslag's theorem.

References

[Bau65] Gilbert Baumslag, *Residual nilpotence and relations in free groups*, J. Algebra **2** (1965), 271–282. MR MR0179239 (31 #3487)

- [LM09] Larsen Louder and D. B. McReynolds, *Graphs of subgroups of free groups*, *Algebr. Geom. Topol.* **9** (2009), no. 1, 327–335. MR 2482080 (2009m:20035)
- [Lou06] Larsen Louder, *Krull dimension for limit groups III: Scott complexity and adjoining roots to finitely generated groups*, *ArXiv Mathematics e-prints* (2006).
- [LS62] R. C. Lyndon and M. P. Schützenberger, *The equation $a^M = b^N c^P$ in a free group*, *Michigan Math. J.* **9** (1962), 289–298. MR MR0162838 (29 #142)
- [Sch59] Eugene Schenkman, *The equation $a^n b^n = c^n$ in a free group*, *Ann. of Math. (2)* **70** (1959), 562–564. MR MR0104723 (21 #3476)
- [Sco73] G. P. Scott, *Finitely generated 3-manifold groups are finitely presented*, *J. London Math. Soc. (2)* **6** (1973), 437–440. MR MR0380763 (52 #1660)
- [She55] Abe Shenitzer, *Decomposition of a group with a single defining relation into a free product*, *Proc. Amer. Math. Soc.* **6** (1955), 273–279. MR MR0069174 (16,995c)
- [Sta65] John R. Stallings, *A topological proof of Grushko’s theorem on free products*, *Math. Z.* **90** (1965), 1–8. MR MR0188284 (32 #5723)
- [Sta83] ———, *Topology of finite graphs*, *Invent. Math.* **71** (1983), no. 3, 551–565. MR MR695906 (85m:05037a)
- [Swa04] G. A. Swarup, *Delzant’s variation on Scott complexity*, January 2004.

Department of Mathematics
 University of Michigan
 Ann Arbor, MI 48109-1043
 USA
email:llouder@umich.edu, lars@d503.net